

WARNING KEEP OUT OF THE REACH OF CHILDREN

PEEL
HERE

IPPON[®] 500SC

FUNGICIDE

Broad spectrum fungicide for the control of specific diseases in a range of crops.

Active Ingredient: Contains 500 g/litre iprodione in the form of a suspension concentrate.

GROUP

2

FUNGICIDE



AGRECOVERY
Rural Recycling Programme

Registered pursuant to the ACVM Act 1997 No. P5644
See www.foodsafety.govt.nz for registration conditions
HSNO Classifications: 6.9B, 9.1A

In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only call: 0800 734 607

NET
CONTENTS:

10

LITRES

AR202/10L



9 421903 469106 >



adria
CROP PROTECTION SOLUTIONS

WARNING KEEP OUT OF THE REACH OF CHILDREN

IPPON[®] 500SC

FUNGICIDE

Broad spectrum fungicide for the control of specific diseases in a range of crops.

Active Ingredient: Contains 500 g/litre iprodione in the form of a suspension concentrate.

GROUP 2 FUNGICIDE



AGRECOVERY
Rural Recycling Programme

Registered pursuant to the ACVM Act 1997 No. P5644
See www.foodsafety.govt.nz for registration conditions
HSNO Classifications: 6.9B, 9.1A

In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only call: 0800 734 607

PHIL STIC MULTIFOLD[®]

NET
CONTENTS:

10

LITRES



AR202/10L



9 421903 469106 >

READ LABEL BEFORE USE. SHAKE WELL BEFORE USE.



HSNO Classifications: 6.9B, 9.1A

Approved under the HSNO Act 1996, HSR000623. See www.epa.govt.nz for controls.

Qualified Person/Contractor: Required.

WARNING: May cause damage to organs through prolonged or repeated exposure.

PRECAUTIONS: Keep out of reach of children. Wear protective clothing such as impervious gloves, waterproof hat, coat and trousers when using. Avoid contact with skin or eyes. Do not breathe fumes, vapours or spray. Wash hands and exposed skin after use and before meals. When mixing or applying wear waterproof gloves. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area. Do not apply directly into or onto water.

STORAGE: Store locked up in the original, unopened container in a cool, dry place, out of direct sunlight and away from stockfeed or foodstuffs. Protect against freezing. As a Class 9 Substance with Ecotoxicity Classifications, storage must be carried out in such a manner as to prevent contamination of waterways. It is recommended that The New Zealand Standard for the Management of Agrichemicals (NZS8409) is followed. See Safety Data Sheet for further information.

PRODUCT DISPOSAL: Product should be used up according to label directions. If this is not possible, then product should be disposed of at an approved facility.

CONTAINER DISPOSAL: Triple rinse container and add residue to spray tank. Return empty container to an AgRecovery collection point for disposal.

ECOTOXIC - VERY TOXIC TO AQUATIC LIFE WITH LONG-LASTING EFFECTS. Avoid contamination of any water supply with chemical or empty container. Avoid release to the environment. A strategy to minimise spray drift should be employed at all times when applying spray near aquatic environments.
contact: After contact with skin, wash immediately with soap and water.

FIRST AID: If swallowed do NOT induce vomiting. For advice, contact the National Poisons Centre on 0800 POISON (0800 764766) or a doctor immediately. If medical advice is needed, have product container or label at hand. Get medical advice/attention if you feel unwell.

Eye contamination: If product or spray enters eyes, wash out immediately and continue rinsing with clean water for at least 15 minutes. Obtain medical attention.

Skin contact: After contact with skin, wash immediately with soap and water.

RESISTANCE MANAGEMENT: Ippon 500SC is a **GROUP 2 FUNGICIDE**. Resistance to this dicarboximide fungicide and related products could develop from excessive use. To minimise this risk use strictly in accordance with label instructions and resistance management strategies that exist for any fungus listed on the label. Avoid using this fungicide exclusively all season and alternate with fungicides from alternative chemical groups.

Ippon 500SC is a contact fungicide of the dicarboximide group.

It is recommended that when used for botrytis control no more than two applications of Ippon 500SC are made per season and that alternative nondicarboximide products are used at other key times in the programme. Do not use this product if resistance monitoring tests show a high level of tolerant strains are present before application is planned.

SPILLS: Wear protective equipment to prevent skin and eye contamination and inhalation of vapors. Contain and prevent contamination of drains and waterways. Use absorbent, soil, vermiculite, atapulgitte bentonite or clay absorbent. Collect and seal in plastic lined drums for disposal in an authorised landfill. Wash the spill zone down with excess water avoiding contamination of any water body.

DIRECTIONS FOR USE

Partly fill the spray tank, add the required amount of Ippon 500SC and top up with water. Agitate thoroughly before spraying, and avoid over-vigorous agitation at any stage. Ippon 500SC is unstable in alkaline solutions: if alkaline water is used to prepare spray solutions use an acidifying or buffering agent to reduce the pH of the final solution to below pH 7. Ippon 500SC is compatible with most commonly used fungicides and insecticides. Do not mix with strongly alkaline materials.

IT IS AN OFFENCE FOR USERS OF THIS PRODUCT TO CAUSE RESIDUES EXCEEDING THE RELEVANT MRL IN THE FOOD NOTICE:

MAXIMUM RESIDUE LEVELS FOR AGRICULTURAL COMPOUNDS.

IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS.

RECOMMENDATIONS

NOTE: The following recommendations are for high volume spraying to the point of run off. For concentrate spraying, adjust dilution rates accordingly. Do not use less than 75ml Ippon 500SC/100 litres (1.5 litres/hectare).

CROP	DISEASE	RATE	WITHHOLDING PERIOD	REMARKS
Grapes	Grey Mould (Botrytis)	75-100ml/100 litres	1 day	Apply once or twice in the period from veraison to harvest using alternative, non-dicarboximides for the rest of the season.
Stonefruit	Brown Rot (Monilinia spp)	75ml/100 litres	1 day	Apply 2-3 sprays at 7-10 day intervals (more frequently following heavy rainfall) in the four week period prior to harvest. Use alternative nondicarboximide fungicides for the rest of the season.
Kiwifruit	Botrytis & Sclerotinia Fruit Rots	75ml/100 litres	1 day (24 hours)	Sclerotinia control: Apply at flowering and again in January if required. Botrytis control: To reduce losses from botrytis storage rot, apply one day before harvest. Note: Applications at flowering for sclerotinia control will also suppress botrytis. DO NOT make more than two applications in any one season.
Tangelos	Alternaria leaf spot (Alternaria citri)	100ml/100 litres	14 days	To reduce losses from Alternaria citri: Apply when last seasons crop has been harvested completely. Apply in a 21-28 day programme with alternative products from the start of flowering. Use the shorter interval during periods of high disease pressure.
Strawberries and other Berryfruit**	Botrytis	100ml/100 litres	1 day	Apply in a programme with alternative products on a 7-10 day schedule from the start of flowering. Reserve use for when infection pressure is high, particularly in the pre-harvest period.
Asparagus	Stemphylium	1.5 litres/ hectare	-	Apply between close up and the end of February at first appearance of the disease on a 21-28 day schedule. Use the shorter interval during periods of prolonged wet weather.

CROP	DISEASE	RATE	WITHHOLDING PERIOD	REMARKS
Fine Turf & Cotula	Fusarium, Red Thread, Brown Patch, Dollar Spot, Helminthosporium Leaf Blight, Curvularia Leaf Blight	50-100*ml in 5 litres water per 100 square metres	-	Apply at 14-21 day intervals in a programme with alternative products from the onset of disease or whilst conditions favour disease.

*Use the higher rate under conditions of severe disease pressure.

**Do not use on Rabbiteye blueberry cultivars.

Notice to Buyer: Careful tests have proved that the product is suitable for the recommended purposes when used in accordance with our instructions. However, since storage and use of product are beyond our control, we do not therefore accept any liability for damage or loss of efficacy which may result from the use of this product, whether used in accordance with the directions or not. We are liable for a consistent quality of the product, but the risk of its storage and use is not borne by us.

Registered to and Distributed by:

Adria New Zealand Limited.

P.O. Box 535, 407 State Highway 16 Kumeu, Auckland.

Ph: 09 412 9817. Fax: 09 412 9807. www.adria.nz

Batch number and manufacturing date: See on packaging

Manufactured by: Chimac sprl, Rue de Renory 26/1, B-4102 Ougree, Belgium.

©Ippon 500SC is a registered trademark of Arysta LifeScience Benelux sprl

Proper Shipping Name	Hazard Class	UN No.	Hazchem	Packing Group
ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (IPRODIONE 50% SOLUTION),	9	3082	3Z	III
Do not carry this product on a passenger service vehicle.				