

## Corona<sup>®</sup> 250WG

A systemic fungicide for the control of Brownrot in stonefruit



Active Ingredient: 250g/kg tebuconazole

**Chemical Group:** DMI-Triazole

Formulation: Water dispersible granule (WG)

Pack Size: 1 kg

## **General Information:**

Corona 250WG is a proven DMI fungicide for the control of Brownrot in stonefruit. It provides excellent efficacy when used in a managed programme with alternative chemical groups. The WG formulation offers tankmixing flexibility and easy mixing characteristics.

- ✓ Protection and curative activity
- √ Good PHI flexibility for local export markets
- √ High quality WG formulation
- ✓ Excellent crop safety

## **Directions for use**

Apply as a dilute spray, ensuring complete coverage, without excessive run-off. For concentrate spraying, adjust dilution rate accordingly.

Crop	Disease	Rate	Critical Comments
Summerfruit	Brownrot	40g/100 litres of water. Use minimum of 800g/ha.	A: Apply in a regular blossom programme from pink to shuck fall inclusive. Thereafter apply alternative non-DMI fungicides through until harvest. or B: Apply 2-3 sprays at 7-10 day intervals in the period prior to harvest. When used pre-harvest use alternative non-DMI fungicides earlier in the season

Apply a maximum of 3 applications, preferably with non-DMI fungicides, as required over the pink to shuck fall period. Thereafter apply alternative non-DMI fungicides through until harvest.

## Directions for use cont...

Mixing: Add the required amount of Corona 250WG into the half-filled spray tank with agitation operating. Continue agitation during spraying and after a stoppage before recommencing spraying.

**Application:** Make sure that spray equipment is correctly calibrated. Ensure complete spray coverage for best results. Do not overspray.

**Compatibility:** Corona 250WG is compatible with most of the commonly used insecticides and fungicides, except alkaline materials such as Bordeaux mixture and lime sulphur.

Withholding Period: Summerfruit -1 day

Resistance Management: Resistance to Corona 250WG and related DMI fungicides could develop from repeated use. To minimise this risk, use strictly in accordance with label instructions. DO NOT make more than 3 applications during the season.