

# WARNING

KEEP OUT OF REACH OF CHILDREN

# Probait

## Snail and Slug Bait

All weather slug and snail bait.

**Active Ingredient: contains 15g/kg metaldehyde  
in the form of a bait.**

Contains deterrent to help with animal safety.

Registered pursuant to the ACVM Act 1997 No. P9358.  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

In a transport emergency dial 111, Police or Fire Brigade.  
For specialist advice in a First Aid emergency only call:  
National Poisons Centre on 0800 764 766.



9 421903 419699 >

**Net Contents: 15kg**



## READ LABEL BEFORE USE



Approved under the HSNO Act 1996, HSR000142. See [www.epa.govt.nz](http://www.epa.govt.nz) for controls.  
HSNO Classifications: 6.8B, 6.9B, 9.1D

**WARNING:** Suspected of damaging fertility or the unborn child. May cause damage to organs through prolonged or repeated exposure. May cause long lasting harmful effects to aquatic life.

**ECOTOXIC: Harmful to the aquatic environment with long lasting effects.**

Avoid contamination of any water supply with chemical or empty container. Must not be applied onto or into water. Harmful to terrestrial vertebrates. Keep away from animals, poultry, ducks and domestic pets as accidental poisoning may be fatal. If pets or livestock are poisoned contact a veterinary surgeon.

**SAFETY DIRECTIONS:**

Keep out of reach of children. Read label before use. Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Do not breathe fumes, vapours and spray. Avoid skin and eye contact. When handling product, wear protective clothing such as waterproof gloves, eye protection, overalls and boots. Remove protective clothing and wash hands, face and exposed skin thoroughly before meals and after work. Do not eat, drink or smoke while using this product. Avoid release to the environment. Use personal protective equipment as required. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area.

**STORAGE:**

Store locked up in the original, unopened container in a cool, dry place, out of direct sunlight and away from stockfeed or foodstuffs. As a Class 9 Substance with Ecotoxicity Classifications, storage must be carried out in such a manner as to prevent contamination of waterways. It is recommended that The New Zealand Standard for the Management of Agrichemicals (NZS8409) is followed. See Safety Data Sheet for further information.

**EQUIPMENT:**

Apply using well maintained and accurately calibrated equipment. Flush equipment thoroughly with clean water after use.

**DISPOSAL:**

Ensure container is completely empty by emptying into application equipment or other use. Burn empty container in an appropriate incinerator if permitted and circumstances, such as wind direction permit, otherwise bury in a suitable landfill. Dispose of unused product only by using in accordance with label directions or at an approved landfill.



**FIRST AID:** If swallowed do NOT induce vomiting. For advice, contact the National Poisons Centre on 0800 POISON (0800 764766) or a doctor immediately. Get medical advice/attention if you feel unwell. IF exposed or concerned: Get medical advice/ attention.

**Batch No. and Manufacturing Date:** See on packaging  
**Registered to and Distributed by:** Adria New Zealand Limited.  
P.O. Box 535, 407 State Highway 16 Kumeu, Auckland.  
Ph: 09 412 9817. Fax: 09 412 9807. [www.adria.nz](http://www.adria.nz)



**Notice to Buyer:** Careful tests have proved that the product is suitable for the recommended purposes when used in accordance with our instructions. However, since storage and use of product are beyond our control, we do not therefore accept any liability for damage or loss of efficacy which may result from the use of this product, whether used in accordance with the directions or not. We are liable for a consistent quality of the product, but the risk of its storage and use is not borne by us.

**PRODUCT INFORMATION:** Pro bait Snail and Slug bait kills slugs and snails by irreversibly destroying the mucus producing cells that are essential for movement.

Where the attack is heavy, it may be necessary to repeat the application. Best results are achieved by application during mild damp weather when slugs and snails are most active but when rain is not imminent.

**DIRECTIONS FOR USE:** Scatter baits evenly using an appropriate applicator by ground or direct drill with seed for slug control. DO NOT place in heaps. Clean up any spilled baits, including any from after filling the applicator.

**TIMING:** Apply baits when crops are most vulnerable. For broadacre crops this is likely to be at or immediately after drilling and emergence. For other crops, time the application to protect the sensitive growth stages. For seedlings best results may be achieved by applying 5-10 days prior to planting. See your distributor for further advice. DO NOT apply to edible plant parts or to crops where baits may be collected with harvested commodities or in animal fodder. DO NOT heap baits.

**RECOMMENDATIONS:** Use higher rates where infestation levels are high or under heavy rainfall conditions. By law, the maximum application rate must not be exceeded.

**Broadcast:** Use 12-18kg/ha (120-180g/100m<sup>2</sup>). Use the high rate where slug/snail levels are high. Broadcast evenly over the surface of the ground or band along the plant/drill rows. Apply by hand or through most granule or fertiliser applicators. For crop establishment apply baits prior to sowing (1-5 days), at sowing, or immediately after sowing. For emerged or established crops apply immediately damage occurs.

**Drilling with seed:** Drill 12kg/ha into the furrow with the seed. Use only in situations where the drill leaves a distinct open drill slot allowing the slugs to easily access to the germinating seeds in the furrow (if in doubt broadcast baits as above). Distribute through granule applicator hopper or mix thoroughly with the seed.

**IT IS AN OFFENCE FOR USERS OF THIS PRODUCT TO CAUSE RESIDUES EXCEEDING THE RELEVANT MRL IN THE FOOD NOTICE: MAXIMUM RESIDUE LEVELS FOR AGRICULTURAL COMPOUNDS.**

**WITHHOLDING PERIOD:**

Do not apply to plant parts for human consumption or plant parts that will be grazed or fed to animals. Withhold stock from treated areas for 7 days.