

Vandia® 250EC

A systemic fungicide for control of stripe rust, leaf rust, powdery mildew and speckled leaf blotch in wheat, scald and leaf rust in barley, crown rust in oats, rusts in ryegrass seed crops, powdery mildew in peas and white rot in onions.



Active Ingredient: 250g/litre triadimenol
Chemical Group: Triazole – GROUP 3 FUNGICIDE
Formulation: Emulsifiable concentrate (EC)
Pack Size: 5 Litre

Vandia® offers highly effective disease control:

- ✓ Systemic activity.
- ✓ Residual control.
- ✓ Broad spectrum.

About Vandia 250EC®

Vandia® is a systemic fungicide, with proven efficacy against white rot in onions, as well as diseases of other crops. It offers both systemic and residual activity to protect the crop from infection.

General Information:

Mixing: Half fill the spray tank with water then pour in the required quantity of Vandia with agitators running, then top up with water. Do not pre-mix.

NOTE: Resistance to this fungicide and related DMI products could develop in some disease situations from repeated use. To minimise this risk use strictly in accordance with label instructions.

IT IS AN OFFENCE for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

Recommendations:

Crop	Disease	Rate	Remarks
Wheat	Stripe rust, Leaf rust, Powdery mildew, Speckled leaf blotch	500ml/ha	Apply on first appearance of disease. Should disease pressure persist a further application may be required 3-4 weeks later.
Oats	Crown rust	500ml/ha	Apply on first appearance of disease. Should disease pressure persist a further application may be required 3-4 weeks later.
Barley	Scald, Leaf rust	500ml/ha	Apply on first appearance on disease.

Resistance Management: In cereals use a maximum of three DMI applications, including seed treatment, per crop, at full recommended label rates.

Crop	Disease	Rate	Remarks
Ryegrass seed crops	Crown rust, Stem rust	500ml/ha	Apply on first appearance of disease. Should disease pressure persist a further application may be required 3-4 weeks later.
Peas	Powdery mildew	250ml/ha	Apply preventatively at flowering or at first sign of disease. A second application may be required 14 days later.

Apply Vandia in at least 200 litres of water per hectare by ground application or 50 litres per hectare by aerial application. Use higher water volumes in dense or advanced crops, especially under dry weather conditions.

Crop	Disease	Rate	Remarks
Onions	White rot	1.5 litres/ha	Apply in 1000 litres of water/ha. Make first application 6-8 weeks after sowing and follow with two further applications at 3-4 week intervals. The use of a suitable seed treatment is recommended.

Compatibility: Vandia is compatible with commonly used insecticides and fungicides.

Withholding Periods: Cereals 49 days, Ryegrass seed crops 35 days, Peas 28 days, Onions 35 days.