

DIRECTIONS FOR USE

It is an offence to use this product on animals.

Apply in a minimum of 150 litres of water per hectare by ground or 60 litres of water per hectare by air.

Apply no more than two Pyrax treatments per season.

Crop	Disease	Rate	Comments
Barley	Scald, Net blotch (spot & net form), Leaf rust, Ramularia leaf and awn spot, Spot blotch, Powdery mildew.	0.8 litre/ha Pyrax plus 0.6 litre/ ha Accuro 125SC .	Apply as a preventative spray on susceptible cultivars and when conditions favour disease. Otherwise, apply at the first sign of disease. Ongoing disease pressure or reinfection may require a further treatment 3-5 weeks later. Apply up to the end of ear emergence (Zadok GS 59).
Wheat	Stripe rust, Leaf rust, Speckled leaf blotch, Glume blotch, Powdery mildew, Sooty moulds, Didymella leaf spot.	0.8 litre/ha Pyrax plus 0.6 litre/ ha Accuro 125SC .	Apply as a preventative spray on susceptible cultivars and when conditions favour disease. Otherwise, apply at the first sign of disease. Ongoing disease pressure or reinfection may require a further treatment 3-5 weeks later. Apply up to the end of ear emergence (Zadok GS 59). Note: Application may be necessary early in the season at mid-tillering to control Speckled leaf blotch.
Ryegrass	Stem rust	0.8 litre/ha Pyrax plus 0.6 litre/ ha Accuro 125SC .	Apply as a preventative spray on susceptible cultivars and when conditions favour disease. Otherwise, apply at the first sign of disease. Ongoing disease pressure or reinfection may require a further treatment 3-5 weeks later. Apply up to the end of flowering (Zadok GS 69).
Maize	Common rust	1.0 litre/ha	Apply as a preventative spray on susceptible cultivars when conditions favour disease, otherwise, and on more tolerant cultivars, apply at the first appearance of disease. Ongoing disease pressure or re-infection may require another spray 14-21 days later. Do NOT apply later than the tasselling stage. Do NOT apply more than 2 applications per season.

Mixing: Add the required amount directly into the partly filled spray tank with agitation operating. Complete filling and maintain agitation until spraying is complete.

Compatibility: Pyrax may be tank mixed with most commonly used insecticides and fungicides, such as the triazole fungicide Accuro 125SC.

Rainfall: Do NOT apply if rainfall is likely within 2 hours of treatment.

Withholding Periods:

IT IS AN OFFENCE FOR USERS OF THIS PRODUCT TO CAUSE RESIDUES EXCEEDING THE RELEVANT MRL IN THE FOOD NOTICE: MAXIMUM RESIDUE LEVELS FOR AGRICULTURAL COMPOUNDS.

Wheat and barley grain: Do NOT harvest within 56 days of application (last application at end of ear emergence - GS 59). Cereal silage/green feed: Allow 28 days between last application and cutting or introduction of stock for grazing.

Maize grain: Do NOT harvest within 56 days of application.

Maize silage/green feed: Allow 28 days between last application and cutting.

Ryegrass seed: Allow 35 days between last application and introduction of stock for grazing.



UN3082. Class: 9, Packaging group: III. ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (PYRACLOSTROBIN 25%).

Do not carry this product on a passenger service vehicle.

DANGER
KEEP OUT OF REACH OF CHILDREN

Pyrax®

Systemic fungicide for the control of diseases
in barley, maize, ryegrass and wheat.

Active Ingredient: contains 250g/litre
pyraclostrobin in the form of an
emulsifiable concentrate.

GROUP 11 FUNGICIDE

Registered pursuant to the ACVM Act 1997 No. P9203
See www.foodsafety.govt.nz for registration conditions

In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only call:
0800 734 607 (24 hours).



Net Contents: 10 litres



READ LABEL BEFORE USE



HSNO Classifications: 3.1D, 6.1D (Inhalation), 6.3A, 6.4A, 6.9A, 9.1A
Approved under the HSNO Act 1996, HSR101115. See www.epa.govt.nz for controls.
Qualified/Competent Person: Required.

DANGER: Combustible liquid. Harmful if inhaled. Causes skin irritation. Causes serious eye irritation. Causes damage to organs through single, prolonged or repeated exposure.

ECOTOXICITY: VERY TOXIC TO AQUATIC LIFE WITH LONG-LASTING EFFECTS. DO NOT contaminate an aquatic environment with chemical or empty container. DO NOT spray over, or allow drift to drift over, an aquatic environment. Must not be applied directly onto or into water. DO NOT allow drift outside the target area to occur.

HANDLING PRECAUTIONS: Read label before use. Keep away from heat, sparks, open flames or hot surfaces. No smoking. Do not breathe fumes, vapours or spray. Avoid contact with eyes and skin. Avoid inhalation of vapour or spray mist. Do not eat, drink or smoke while using. Remove protective clothing and wash hands and face thoroughly before meals and after work. Apply using calibrated and maintained equipment in accordance with the NZ Standard for the Management of Agrichemicals (NZS8409). Use only outdoors or in a well-ventilated area. In case of fire evacuate area.

STORAGE: Store locked up in the original, unopened container in a cool, well-ventilated, dry place, out of direct sunlight and away from stockfeed or foodstuffs. As a Class 9 Substance with Ecotoxicity Classifications, storage must be carried out in such a manner as to prevent contamination of waterways. It is recommended that The New Zealand Standard for the Management of Agrichemicals (NZS8409) is followed. See Safety Data Sheet for further information.

SPRAY DRIFT: The person applying the substance must not cause adverse effects beyond the boundary of the treated property, and must also avoid adverse effects from spray drift occurring. Mitigation measures employed must be recorded as part of the application records.

PERSONAL PROTECTIVE EQUIPMENT (PPE): When mixing or applying, wear chemical resistant overalls, chemical resistant boots and gloves, chemical resistant footwear plus socks, chemical resistant head gear and eye protection. Clean protective clothing daily after work.

APPLICATION RESTRICTIONS:

- Pyrax must not be applied at rates exceeding 1L of formulated product/ha per application (equivalent to 250g pyraclostrobin/ha); and
- Pyrax must not be applied more than twice per crop cycle per 365 days; and
- An interval of at least 14 full days must be observed before Pyrax is reapplied to the same area.

SPILLAGE: Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely, according to local Council regulations.

DISPOSAL: Triple rinse empty container and add rinsate to spray tank. Triple rinsed containers should be taken to an AgRecovery collection site. Otherwise crush and bury in a suitable landfill. Do not burn empty containers.



First Aid: If medical advice is needed, have product container or label at hand. IF ON SKIN: Wash with plenty of soap and water. Take off contaminated clothing and wash before re-use. If skin irritation occurs: Get medical advice/ attention. IF INHALED: Remove to fresh air and keep at rest in a position comfortable for breathing. Call a POISON CENTER or doctor/physician if you feel unwell. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention. If exposed: Call a POISON CENTER or doctor/physician. In case of fire: Use dry chemical extinguisher, foam or carbon dioxide to extinguish.

Call 09 412 9817 or visit: www.adria.nz

Batch No. and Manufacturing Date: See on packaging

Registered to and Distributed by: Adria New Zealand Ltd.

P.O. Box 5354

407 State Highway 16, Kumeu, Auckland

Ph: 09 412 9817 Fax: 09 412 9807

www.adria.nz



Notice to Buyer: Careful tests have proved that the product is suitable for the recommended purposes when used in accordance with our instructions. However, since storage and use of product are beyond our control, we do not therefore accept any liability for damage or loss of efficacy which may result from the use of this product, whether used in accordance with the directions or not. We are liable for a consistent quality of the product, but the risk of its storage and use is not borne by us.

GENERAL INFORMATION: Pyrax is a strobilurin fungicide with eradicant and protectant activity. It also has translaminar activity which provides some protection to new growth. For best results, thorough coverage of the crop is essential. Optimal results are achieved with application before the onset of infection.

RESISTANCE MANAGEMENT: Pyrax is a **GROUP 11 FUNGICIDE**. Resistance to this fungicide and related QoI products could develop in some disease situations from repeated use. To minimise this risk, use strictly in accordance with the label instructions and resistance management strategies that exist for any fungus listed on the label. Always apply Pyrax in a tank mix as directed or with a registered fungicide from a different mode of action group. Follow individual crop resistance management strategies. Refer to the NZCPR website for more detailed information (<https://resistance.nzpps.org/>).