

GENERAL INFORMATION: Mixing: Half fill the spray tank with water then pour in the required quantity of Vandia 250EC with agitators running, then top up with water. Do not pre-mix.

NOTE: Vandia is a GROUP 3 FUNGICIDE from the DMI/Triazole group of fungicides. Resistance to this fungicide and related DMI products could develop in some disease situations from repeated use. To minimise this risk use strictly in accordance with label instructions. Apply no more than 3 DMI applications in a season.

IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS.

Recommendations:

Crop	Disease	Rate	Remarks
Wheat	Stripe rust, Leaf rust, Powdery mildew, Speckled leaf blotch	500ml/ha	Apply on first appearance of disease. Should disease pressure persist a further application may be required 3-4 weeks later.
Oats	Crown rust	500ml/ha	Apply on first appearance of disease. Should disease pressure persist a further application may be required 3-4 weeks later.
Barley	Scald, Leaf rust	500ml/ha	Apply on first appearance on disease.

Resistance Management: In cereals use a maximum of three DMI applications, including seed treatment, per crop, at full recommended label rates.

Ryegrass seed crops	Crown rust, Stem rust	500ml/ha	Apply on first appearance of disease. Should disease pressure persist a further application may be required 3-4 weeks later.
Peas	Powdery mildew	250ml/ha	Apply preventatively at flowering or at first sign of disease. A second application may be required 14 days later.

Apply Vandia 250EC in at least 200 litres of water per hectare by ground application or 50 litres per hectare by aerial application. Use higher water volumes in dense or advanced crops, especially under dry weather conditions.

Onions	White rot	1.5 litres/ha	Apply in 1000 litres of water/ha. Make first application 6-8 weeks after sowing and follow with two further applications at 3-4 week intervals. The use of a suitable seed treatment is recommended.
---------------	-----------	---------------	--

COMPATIBILITY: Vandia 250EC is compatible with commonly used insecticides and fungicides.

IT IS AN OFFENCE for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS:

Cereals 49 days, Ryegrass seed crops 35 days, Peas 28 days, Onions 35 days



UN3082. CLASS 9, PG III. ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (TRIADIMENOL 25%). Do not carry this product on a passenger service vehicle.

DANGER
KEEP OUT OF REACH OF CHILDREN

Vandia® 250EC

A systemic fungicide for control of stripe rust, leaf rust, powdery mildew and speckled leaf blotch in wheat, scald and leaf rust in barley, crown rust in oats, rusts in ryegrass seed crops, powdery mildew in peas and white rot in onions.

Active Ingredient: contains 250g/litre triadimenol in the form of an emulsifiable concentrate. Also contains 1-Methyl 2-pyrrolidinone.

Group 3 Fungicide

Registered pursuant to the ACVM Act 1997 No. P8412
See www.foodsafety.govt.nz for registration conditions.

In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only call: 0800 734 607 (24 hours).



Net Contents: 5 litres



READ THIS LABEL COMPLETELY BEFORE USE



HSNO Classifications: 6.1D (Oral), 6.3A, 6.4A, 6.8A, 6.9A, 9.1B, 9.1C, 9.2C, 9.3C
Approved under the HSNO Act 1996, HSR000082. See www.epa.govt.nz for controls.

DANGER: Harmful if swallowed. Causes skin irritation. Causes serious eye irritation. May damage fertility or the unborn child. Causes damage to organs through prolonged or repeated exposure. Toxic to aquatic life with long lasting effects. Harmful to the soil environment. Harmful to terrestrial vertebrates.

SAFETY DIRECTIONS: Keep out of reach of children. Read label before use. Do not handle until all safety precautions have been read and understood. Do not breathe fumes, vapours or spray. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Avoid release to the environment. Use personal protective equipment as required. Collect Spillage. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area. Do not apply directly into or onto water.

STORAGE: Store locked up in the original, unopened container in a cool, dry place, out of direct sunlight and away from stockfeed or foodstuffs. As a Class 9 Substance with Ecotoxicity Classifications, storage must be carried out in such a manner as to prevent contamination of waterways. It is recommended that The New Zealand Standard for the Management of Agrichemicals (NZS8409) is followed. See Safety Data Sheet for further information.

CONTAINER DISPOSAL: Triple rinse container and add residue to spray tank. Return empty container to an AgRecovery collection point for disposal.

TOXIC TO AQUATIC ORGANISMS. Avoid contamination of any water supply with chemical or empty container. A strategy to minimise spray drift should be employed at all times when applying spray near aquatic environments.



FIRST AID: If medical advice is needed, have product container or label at hand. Get medical advice/attention if you feel unwell. IF SWALLOWED: Rinse mouth. Call a POISON CENTER or doctor/physician if you feel unwell. IF ON SKIN: Wash with plenty of soap and water. If skin irritation occurs: Get medical advice/ attention. Take off contaminated clothing and wash before re-use. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention. IF exposed or concerned: Get medical advice/ attention.

SPILLS: Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely, according to local Council regulations.

Notice to Buyer: Careful tests have proved that the product is suitable for the recommended purposes when used in accordance with our instructions. However, since storage and use of product are beyond our control, we do not therefore accept any liability for damage or loss of efficacy which may result from the use of this product, whether used in accordance with the directions or not. We are liable for a consistent quality of the product, but the risk of its storage and use is not borne by us.

Registered to and Distributed by:

Adria New Zealand Limited.
P.O. Box 535, 407 State Highway 16 Kumeu, Auckland.
Ph: 09 412 9817. www.adria.nz



Batch number and manufacturing date: See on packaging