

**DIRECTIONS FOR USE:**

**Mixing:** Shake well before use. Add the required amount to a partly filled spray tank with agitation system operating. Complete filling the tank with water and apply. Flush spray system with clean water after use.

**Compatibility:**

DO NOT USE CORONET IN ANY MIXTURES WHEN APPLYING TO PERENNIAL RYEGRASS.

When used on wheat, Coronet can be tankmixed with Accuro 125SC and chlormequat-chloride in 2 and 3 way mixtures. Care must be taken to ensure that recommended timings for all products are followed. The addition of a suitable adjuvant oil at label rates is recommended. Do not use Coronet within 5 days of broad-leaved herbicide applications.

CROP	WEEDS	RATE	REMARKS
Wheat	Wild oats	750mL/ha in 250 litres of water. Add adjuvant oil at recommended rates.	Apply when wild oats are between 3 leaf and 1st node visible growth stage and when they are growing actively.
	Lesser canary grass ( <i>Phalaris minor</i> )	750mL/ha in 250 litres of water. Add adjuvant oil at recommended rates.	Apply when weeds are between the 3 leaf visible stage and the end of tillering and are growing actively. Avoid application during periods of moisture stress.
Perennial ryegrass ( <i>Lolium perenne</i> )	Wild oats	750mL/ha in 250 litres of water. <b>Do not use with any other agrochemicals or adjuvants.</b>	Apply in early spring when wild oats are between 3 leaf and end of tillering growth stage and are actively growing. Note: Seed yield response is likely to be reduced the longer the delay in the removal of wild oats in spring.

**DO NOT USE ON BARLEY****IMPORTANT NOTES:**

Gnawed canary grass (*Phalaris paradoxa*) is not controlled by Coronet at label rates.

Do not overlap swath width in ryegrass crops as yield suppression may occur.

Do not use Coronet with any other agrochemical or spray additive when applied to Perennial ryegrass.

Resistant weeds: Annual and Hybrid ryegrasses, tall fescue, chewing's fescue, cocksfoot, browntop, grazing brome (=spiky brome – *Bromus stamineus*), prairie grass and most broadleaved weeds.

Rainfastness: Coronet is rainfast within one hour of application.

**IT IS AN OFFENCE for users of this product to cause residues exceeding the relevant MRL in the Food Notice:**

**Maximum Residue Levels for Agricultural Compounds.**

**Withholding periods:**

Wheat - 70 days (Do not graze treated crop for at least 70 days after treatment)

Perennial ryegrass - 42 days (Do not graze treated crop for at least 42 days after treatment)

**Soil Activity:** Coronet is rapidly broken down by soil microbes causing no restrictions on crop rotations.

Undersown crops: Coronet can be used on crops undersown with certain grasses (See under "Resistant Weeds" above) and clover.

Do not use on crops undersown with "Concord" ryegrass.



**UN 3082. Class: 9, Packaging group: III. ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S.**  
(69g/L fenoxaprop-P-ethyl, 18g/L mefenpyr-diethyl). Do not carry this product on a passenger service vehicle.

# WARNING

KEEP OUT OF REACH OF CHILDREN

# Coronet®

A selective grass herbicide for use on autumn and spring sown wheat and perennial ryegrass for the control of wild oats and lesser canary grass (*Phalaris minor*).

Active Ingredient: contains 69 g/litre fenoxaprop-P-ethyl in the form of an oil-in-water emulsion.

**GROUP A HERBICIDE**

Registered pursuant to the ACVM Act 1997 No. P8196  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions

In a transport emergency dial 111, Police or Fire Brigade.  
For specialist advice in an emergency only call:  
0800 734 607 (24 hours).



**Net Contents: 10 litres**



Approved under the HSNO Act 1996, HSR100440. See [www.epa.govt.nz](http://www.epa.govt.nz) for controls.

HSNO Classifications: 6.3B, 6.5B, 6.9B, 9.1A. Qualified/Competent Person: Required.

This substance may be harmful if swallowed, inhaled or absorbed through the skin, and may cause irritation to eyes and skin. May cause an allergic skin reaction. Do not breathe fumes, vapours or spray. May cause damage to organs through prolonged or repeated exposure.

**PRECAUTIONS:** Avoid contact with skin and eyes, and inhalation of spray mist. Do not eat, drink or smoke while using this product. Wear protective gloves, protective clothing, eye protection and face protection. Wash hands and exposed skin before meals and after work. Contaminated clothing should not leave the workplace and must be cleaned before re-use. Avoid contamination of any water supply with chemical or empty container. Do not apply directly into or onto water. Apply this product carefully. Spray drift may cause serious damage to desirable plants. Ensure compliance with relevant regional air management plans.

**STORAGE:** Store locked up in the original, unopened container in a cool, dry place, out of direct sunlight and away from stockfeed or foodstuffs. As a Class 9 Substance with Ecotoxicity Classifications, storage must be carried out in such a manner as to prevent contamination of waterways. It is recommended that The New Zealand Standard for the Management of Agrichemicals (NZS8409) is followed. See Safety Data Sheet for further information.

**VERY TOXIC TO AQUATIC ORGANISMS WITH LONGLASTING EFFECTS.** A strategy to minimise spray drift should be employed at all times when applying spray near aquatic environments.

**DISPOSAL OF CONTAINER:** Triple rinse container and add residue to spray tank. Return clean, empty container to an AgRecovery site for disposal. Avoid contamination of any water supply with chemical or empty container.



**FIRST AID: Ingestion:** Do NOT induce vomiting if swallowed. Wash mouth with water and seek medical attention immediately. Small sips of water may be given. For advice, contact the National Poisons Centre on 0800 POISON (0800 764766). **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Seek medical attention immediately. **Skin Contact:** Immediately flush body and clothes with large amounts of water. Remove contaminated clothing and footwear. Gently wash affected areas with soap and water. If a large area is affected or if irritation or rash persists seek medical assistance. **Inhalation:** Should irritation occur, remove patient to fresh air. If breathing is shallow or has stopped, ensure airway is clear and apply resuscitation. Seek medical assistance soon if irritation persists. If medical advice is needed, have product container or label at hand. Get medical advice/attention if you feel unwell.

**SPILLS:** Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely (such as to a suitable landfill).

**Batch No. and Manufacturing Date:** See on packaging

**Registered to and Distributed by:** Adria New Zealand Limited.



P.O. Box 535, 407 State Highway 16 Kumeu, Auckland.

Ph: 09 412 9817. Fax: 09 412 9807. [www.adria.nz](http://www.adria.nz)

**Notice to Buyer:** Careful tests have proved that the product is suitable for the recommended purposes when used in accordance with our instructions. However, since storage and use of product are beyond our control, we do not therefore accept any liability for damage or loss of efficacy which may result from the use of this product, whether used in accordance with the directions or not. We are liable for a consistent quality of the product, but the risk of its storage and use is not borne by us.

**Resistant Weeds Warning:** Coronet is a **GROUP A HERBICIDE** from the aryloxyphenoxypropionates (FOPs) chemical group. Naturally occurring weed biotypes resistant to aryloxyphenoxypropionate herbicides can become more prevalent after many years of continuous use. To delay the onset of, or control resistant weeds, use in rotation with herbicides of a different mode of action. Since the occurrence of resistant weeds is difficult to detect prior to use, Adria New Zealand Limited accepts no liability for losses that may result from the failure of this product to control resistant weeds.