

Genoxy® 240sc

A targeted insecticide for controlling leafrollers in grapes and avocados.



Active Ingredient: 240g/litre methoxyfenozide

Chemical Group: Diacylhydrazine

Formulation: Suspension concentrate

Pack Size: 5 litre

Genoxy® offers dependable caterpillar control:

- ✓ Ovi-larvicidal activity controls eggs and caterpillars.
- √ Highly selective to beneficials.
- ✓ Long-lasting activity up to 3 weeks residual activity.
- ✓ No cross-resistance to most other insecticide groups.
- ✓ Low toxicity

About Genoxy® 240SC

Genoxy® is a new generation Moult Accelerating Compound (MAC) insecticide which causes premature moulting in caterpillars. Feeding damage stops within a day, although death may take several days. Genoxy® increases ovicidal activity and will control later caterpillar instars.

Genoxy® can be used in Integrated Pest Management (IPM) programmes and can be applied over flowering where appropriate under the directions for use.

IT IS AN OFFENCE FOR USERS OF THIS PRODUCT TO CAUSE RESIDUES EXCEEDING THE RELEVANT MRL IN THE FOOD NOTICE: MAXIMUM RESIDUE LEVELS FOR AGRICULTURAL COMPOUNDS

Directions for use:

Monitor treated areas for all pests and consult advisers for recommendations to control non-target species. Genoxy is a Moult Accelerating Compound (MAC) insecticide which induces premature moulting in caterpillars. Damage stops within 24 hours but insect death may take a few days.

Genoxy can be utilised in Integrated Pest Management (IPM) programmes and can be used during flowering when appropriate under the directions for use.

The rates suggested are for high volume spraying to run-off. For concentrate spraying adjust dilution rate accordingly. Thorough coverage of plants is essential for control of pests.

Directions for use cont...

CROP	PEST	APPLICATION RATE	APPLICATION TIMING
Avocado	Leafrollers	Apply 25ml per 100 litres to full coverage (point of runoff) with a minimum dose of 500ml/ha. For mature trees and water rates greater than 2000L/ha, maintain the rates of 25ml per 100 litres.	Apply at 3 to 4 week intervals from flowering as part of a season long leafroller spray programme. Do not apply more than four applications in any season.
Grapes		Apply 15-25ml/100L to full coverage. Do not apply Genoxy at less than 150ml/ha under moderate pest pressure and 250ml/ha under severe pest pressure.	Make a single spray application at 80% capfall. If pest pressure remains high at pre-bunch closure, apply an alternative product.

Withholding Periods:

Avocados 14 days (Consult your exporter for WHPs for export markets).

Grapes Do not apply after 80% capfall.

Grazing Restriction: The following restrictions on the use of this product must be observed. DO NOT feed treated crops or pasture to stock other than sheep used for leaf plucking in vineyards. Take care not to allow this product to drift onto pasture or crops that may be grazed. DO NOT graze stock in orchards which have been treated with Genoxy 240SC.

Vineyards: Do not graze stock other than sheep in vineyards which have been treated with Genoxy 240SC. Sheep may be grazed in vineyards for the purpose of leaf plucking providing they are not slaughtered for 2 months after leaving the vineyard.