

Pilan[®] 500SC

Insect growth regulator for the control of mealy bug, scales and whitefly.



Active Ingredient: 500g/litre buprofezin

Chemical Group: Thiadiazine

Formulation: Suspension Concentrate

Pack Size: 5 litre

- ✓ Highly selective IGR.
- ✓ Residual activity.
- ✓ Safe to a range of beneficials - including bumblebees, parasitoid wasps and Encarsia spp.
- ✓ IPM / IWP compatible .
- ✓ Low toxicity.

About Pilan 500SC:

Pilan is an insect growth regulator which has activity against scales, mealy bugs and whitefly. It controls nymphal stages through the moulting process. It may suppress oviposition of adult females and reduce the viability of eggs. Pilan is highly selective to most families of beneficial insects, including the parasitoid wasps. Crop monitoring and targeted applications are critical to gain the best results from Pilan.

General Information: Pilan is an insect growth regulator that interferes with the moulting process in the young developmental stages of mealy bug, scale and whitefly. Pilan has no effect on adults. The action of Pilan on pest populations is gradual, causing a decline in the population as each generation is affected. Pilan is suitable for use in IPM programmes as it is safe to most predators and parasites. It is also safe to bees.

Resistance Management: Pilan is from a unique chemical group. However, overuse may lead to resistance in target populations. To minimise whitefly resistance, use no more than 3 applications per crop per season. Following early season use on tamarillos, alternate with other whitefly materials. Pilan may be alternated with, or mixed with conventional insecticides.

Compatibility Pilan is compatible with most commonly used fungicides and insecticides. Do not mix with strongly alkaline materials such as Bordeaux mixture or lime sulphur.

Mixing: Add the required amount of Pilan directly into the partially-filled spray tank with agitator operating. Maintain agitation during application.

IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS.

Recommendations: The rate of 25mL/100 litres is for high volume application to the point of run-off. For concentrate spraying, adjust dilution rates accordingly.

Directions for use:

Application rate: 25mL/100 litres for all crops. Note recommended water rates for crops. Applications in all crops should ensure that not less than 250mL/ha are applied.

Crop	Pest	Remarks
Citrus	Scales	Apply in 2500-3000 litres of water/ha or more if required to obtain complete coverage. Treat when scale first appears and repeat 10-14 days later. Where populations persist, re-apply at intervals not less than one month. For season long programmes, alternate Pilan with other scale materials..
Grapes	Mealy bug	Apply in a minimum of 1000 litres of water (250mL/ha minimum dose rate). Apply the first spray in mixture with 1% spraying oil at the first sign of bud movement. Make a second application without oil 14-21 days later when shoots are 10-20mm long. Do not apply after the start of flowering.
Persimmon	Mealy bug	Apply two applications at 10-14 days between green tip and bloom. Use sufficient water to obtain complete plant coverage.
Tamarillo	Whitefly	Apply after pruning and before fruit set. Re-apply no sooner than 28 day intervals when whitefly adults exceed 20 per leaf. Alternate with other whitefly materials to control adults.

IT IS AN OFFENCE FOR USERS OF THIS PRODUCT TO CAUSE RESIDUES EXCEEDING THE RELEVANT MRL IN THE FOOD NOTICE: MAXIMUM RESIDUE LEVELS FOR AGRICULTURAL COMPOUNDS.

Withholding periods:

Tamarillo – 7 days

Citrus – 14 days

Grapes – Do not apply after the start of flowering

Registered to and distributed by: Adria New Zealand Limited. P.O. Box 535 Kumeu, Auckland 0841, NEW ZEALAND.

Ph: +64-9-412-9817 www.adria.nz.

Registered pursuant to the ACVM Act 1997, No. P8056. Approved under the HSNO Act 1996, HSR100077.

For more information call us on +64-9-412-9817 | www.adria.nz