

Ivento®

Fungicide for the control of Late blight in potatoes and Downy mildew in onions.



Active Ingredient: 400g/litre cyazofamid
Chemical Group: GROUP 21 FUNGICIDE
Formulation: Suspension concentrate (SC)
Pack Size: 5L

- ✓ Highly effective against Late blight.
- ✓ Excellent protection for early season growth.
- ✓ Rainfast in 1 hour.
- ✓ Unique mode of action and important resistance management tool.
- ✓ Effective against all stages of oomycetes fungal development.

About Ivento®

Ivento® is a fungicide with activity against fungi from the class of Oomycetes, particularly against *Phytophthora infestans* of potatoes and Downy mildew of onions. It acts as a protectant and needs to be applied before the disease infection occurs. It is highly active against Late blight in potatoes and can be used in the early season programme to protect expanding leaves in a rapidly growing crop. Leaf penetration is rapid and Ivento® continues to protect expanding leaves. Repeat applications should be made at 7-10 day intervals to protect new leaves. Ivento® is also an effective resistance management tool when used in conjunction with alternative fungicide groups like mancozeb and cymoxanil, for example.



Directions for use:

Crop	Disease	Maximum Rate	Critical Comments
Potatoes	Late blight (Phytophthora infestans)	200ml/ha	<p>Apply as a protectant spray. Do NOT wait for disease to appear. When conditions are conducive to disease development, apply consecutive sprays of Ivento at a minimum of 7-10 day intervals.</p> <p>Use the shorter interval when infection pressure is high. The total number of Ivento applications per treated crop must not exceed 6. When applying Ivento consecutively, apply a maximum of 3 consecutive sprays, and then switch to a fungicide from an alternative chemical group. Ensure thorough coverage of plants.</p>
Onions	Downy mildew (Peronospora destructor)	200ml/ha + adjuvant	<p>Apply as a protectant, preventative spray, before the appearance of disease when conditions for disease infection are conducive. When conditions are in favour of disease development, apply consecutive sprays of Ivento at a minimum of 7-10 day intervals. The total number of Ivento applications per treated crop must not exceed 6. When applying Ivento consecutively, apply a maximum of 3 consecutive sprays, and then switch to a fungicide from an alternative chemical group. Ensure thorough coverage of plants. Add an appropriate adjuvant.</p>

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS:

Potatoes: 7 days

Onions: 14 days

Do NOT graze or feed treated crops to animals.

RESISTANCE MANAGEMENT: Ivento must be applied according to the label directions. Some naturally occurring individual fungi resistant to the product and other similar fungicides (GROUP 21 FUNGICIDE) may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by this product, thus resulting in a reduction in efficacy and possible yield loss. Since the occurrence of resistant fungi is difficult to detect prior to use, Adria NZ Limited accepts no liability for any losses that may result from the failure of this product to control resistant fungi.