

# Ormax<sup>®</sup>

Fungicide for the control of Late blight in potatoes.



**Active Ingredient:** 62.5g/litre fluopicolide + 625g/litre propamocarb as the hydrochloride

**Chemical Group:** GROUP 43 + 28 FUNGICIDE

**Formulation:** Suspension concentrate (SC)

**Pack Size:** 10L

- ✓ Dual mode-of-action fungicide chemistries with systemic and translaminar activity.
- ✓ Alternative chemical groups to existing potato Late blight fungicides.
- ✓ Field trialled by Plant & Food Research in Pukekohe (2024) on 'Agria', 'Red Rascal' and 'Lady Rosetta' potato varieties at standard and 2x label rates, with no phytotoxic effects in any treatment.
- ✓ Strong early curative and lasting preventative canopy protection.
- ✓ Effective on all major Late blight strains.

## About Ormax<sup>®</sup>

Ormax<sup>®</sup> is a new fungicide with systemic and protectant activity for the control of Late blight (*Phytophthora infestans*) in potatoes. Ormax blocks every stage of Late blight life cycle, with its protectant, curative and anti-sporulant activity. Ormax can be positioned through the season, as Propamocarb's systemic and Fluopicolide's translaminar mobility protect the crop in periods of rapid and stable canopy growth.



**The Resistance Management:** Ormax is a **GROUP 28 + 43 FUNGICIDE**, containing fluopicolide and propamocarb. Diseases resistant to this group of fungicides may develop if overused. To minimise the development of resistance, use strictly in accordance with label recommendations. **Do not use more than 3 times per season.**

**Compatibility:** Ormax is compatible with most fungicides and insecticides. **However, the crop safety of all potential tank-mixes, including additives and other pesticides, has not been tested. Before applying any tank-mixture, safety to the target crop should be confirmed.** To test for crop safety, apply the mixture to a small area of the target crop in accordance with the label instructions to make sure that a phytotoxic effect will not occur. Not compatible with strongly alkaline products.

**IT IS AN OFFENCE FOR USERS OF THIS PRODUCT TO CAUSE RESIDUES EXCEEDING THE RELEVANT MRL IN THE FOOD NOTICE: MAXIMUM RESIDUE LEVELS FOR AGRICULTURAL COMPOUNDS.**

**Withholding Periods:**

Potatoes: 14 days

**IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS.**



**DIRECTIONS FOR USE:**

**Mixing:** Add the required amount of Ormax to a partly filled spray tank with the agitation system operating. Add partner products and complete filling with water. Maintain agitation until spraying is complete. Ensure thorough and even spray coverage throughout the crop.

Crop	Disease	Rate / Ha	Remarks
Potatoes:	Late Blight	1.2 – 1.6 litres in 250-500 litres of water per hectare	Apply at 10-14 day intervals. Use the shorter interval when rapid growth and conditions are conducive to disease. For resistance management it is recommended that no more than 3 sprays be applied in one season.