

# Boscalo®

**Fungicide for the control of various diseases in beans, citrus, grapes, onions and stone fruit.**



**Active Ingredient:** 252g/kg boscalid + 128g/kg pyraclostrobin

**Chemical Group:** GROUP 7 + 11 FUNGICIDE

**Formulation:** Water dispersible granule (WG)

**Pack Size:** 2.5kg

- ✓ **Dual mode-of-action fungicide chemistries with protectant and translaminar activity.**
- ✓ **Group 7 + Group 11 fungicide co-formulation.**
- ✓ **Rapid disease control.**
- ✓ **Efficacy against a range of diseases.**

## About Boscalo®

Boscalo® is a dual mode-of-action fungicide with anti-sporulant and translaminar activity against a range of diseases in beans, citrus, grapes, onions and stone fruit.

Boscalo® effectively inhibits mycelial growth and provides fast disease control and superior protection.

**GENERAL INFORMATION:** Boscalo® is a combination of carboximide and strobilurin fungicides, which provide protectant activity against certain diseases in beans, citrus, grapes, onions and stonefruit. For optimal efficacy, good spray coverage is essential, and applications should be made before disease establishment.



## IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS.

**DIRECTIONS FOR USE:** The recommended rates are for high volume spraying to the point of run-off. For concentrate spraying the rate of product applied must be equal to the rate that would be applied per hectare with dilute application.

Crop	Disease	Rate	Remarks
<b>Beans</b>	Botrytis Sclerotinia white mould	1.5kg/ha in a minimum of 300 litres water/ha.	Apply in a protectant programme. Begin prior to infection and continue at 7-14 day intervals as necessary. Apply as a preventative spray and when conditions favour disease. Alternate with an effective fungicide with a different mode of action. Do NOT apply more than 3 times per season.
<b>Citrus</b>	Alternaria brown spot Citrus scab Melanose	50g/100 litres water.	Apply in a protectant programme from flowering up to harvest. Begin prior to infection and continue at 21 to 28 day intervals. Apply as a preventative spray and when conditions favour disease. Do NOT apply more than 2 consecutive sprays. Alternate with an effective fungicide with a different mode of action. Do NOT apply more than 3 times per season.
<b>Grapes</b>	Botrytis (Botrytis cinerea), Powdery mildew (Uncinula necator), Downy mildew (Plasmopara viticola)	150g/100 litres of water, applying no less than 1.5kg in a minimum of 500 litres water/ha.	Apply in a protectant programme from early flowering to pre-harvest as a preventative spray and when conditions favour disease. Ensure bunch zone is adequately treated. Do NOT apply more than 2 sprays per season. Do NOT apply to hybrid varieties without first testing.
<b>Onions</b>	White rot, [Downy mildew (Plasmopora destructor) – suppression only]	1.6kg/ha in 500 litres water per ha.	Apply in sequence after 1-2 triazole fungicides (e.g. Vandia 250EC) applications. Either apply in a block of 2 consecutive applications of Boscalo, or apply in alternation with an effective fungicide with an alternative mode of action. Do NOT apply more than 2 times per season.
<b>Stonefruit</b>	Blossom blight, Botrytis fruit rot, Rust	50g/100 litres water.	Apply from early flowering up to 100% petal fall (prior to shuck fall) at an interval of 5-14 days with no more than 3 applications per season. Apply as a preventative treatment and when conditions favour disease. Use the shorter interval when disease pressure is high. Do NOT apply more than 2 consecutive sprays of Boscalo. Apply in alternation with an effective fungicide from an alternative chemical group.

## IT IS AN OFFENCE FOR USERS OF THIS PRODUCT TO CAUSE RESIDUES EXCEEDING THE RELEVANT MRL IN THE FOOD NOTICE: MAXIMUM RESIDUE LEVELS FOR AGRICULTURAL COMPOUNDS.

**Withholding periods: Beans:** Do NOT apply within 7 days of harvest; **Citrus:** Do NOT apply within 1 day of harvest;

**Grapes:** Do NOT apply within 28 days of harvest; **Onions:** Do NOT apply within 56 days of harvest;

**Stonefruit (excluding cherries):** Do NOT apply after 100% petal fall (prior to shuck fall).

**Note:** For all export crops, refer to the specific exporter guidelines which may differ from the above.

**Do NOT allow stock to graze within treated orchards until after harvest.**

**Leaf plucking in vineyards:** Where leaf plucking with sheep is carried out in vineyards sprayed with this product, a 2 month slaughter withholding period must be observed after the sheep have been removed from the vineyard.

**Application:** May be applied through all conventional, ground-based boom spraying equipment. Maintain agitation during mixing and spray application.

**Compatibility:** Boscalo may be tankmixed with most commonly used insecticides and fungicides.

**Mixing:** Add the required amount directly into a partly filled spray tank with agitation operating. Complete filling and maintain agitation until spraying is complete.

**Rainfall:** Boscalo is rainfast once dried on the crop.

Registered to and distributed by: Adria New Zealand Limited. P.O. Box 535 Kumeu, Auckland 0841, NEW ZEALAND.  
Ph: +64-9-412-9817; [www.adria.nz](http://www.adria.nz). Boscalo is a Registered Trademark and is Registered pursuant to the ACVM Act 1997, Approval No. P10086. EPA Approval Code: HSR101536.

For more information call us on +64-9-412-9817 | [www.adria.nz](http://www.adria.nz)