

Concero®

Systemic fungicide for the control of diseases in cereals



Active Ingredient: 150g/litre prothioconazole + 75g/litre bixafen

Chemical Group: GROUP 3 + 7 FUNGICIDE

Formulation: Emulsifiable concentrate (EC)

Pack Size: 10L

- ✓ **Dual mode-of-action fungicide chemistries with protectant and curative activity.**
- ✓ **Group 3 (DMI) + Group 7 (SDHI) fungicide co-formulation.**
- ✓ **Rapid disease control.**
- ✓ **Efficacy against a wide range of diseases.**

About Concero®

Concero® is a leading edge fungicide which provides protective and curative activity against a range of diseases in cereals. Combining proven DMI and SDHI chemistries, Concero® offers highly cost-effective disease control with proven yield results and return on investment.



IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS.

DIRECTIONS FOR USE:

CROP	DISEASE	RATE	REMARKS
Wheat	Stripe rust Leaf rust Speckled leaf blotch Tan spot	700-1000ml/ha	Apply at first appearance of disease. Should disease pressure persist or re-infection occur, another application may be required 3-4 weeks later. Use the higher rate when disease pressure is high.
Barley	Scald Net blotch Ramularia leaf spot Leaf rust	700-1000ml/ha	Apply on first appearance of disease. Should disease pressure persist or re-infection occur, another application may be required 3-4 weeks later. Use the higher rate when disease pressure is high.
Triticale	Stripe rust Leaf rust	700-1000ml/ha	Apply on first appearance of disease. Should disease pressure persist or re-infection occur, another application may be required 3-4 weeks later. Use the higher rate when disease pressure is high.

Concero® must only be applied via ground-based methods.

Apply Concero® in at least 120 litres of water per hectare by ground application. Use high water volumes in dense or advanced crops especially under dry weather conditions.

Mixing: Add the required amount directly into partly filled spray tank with agitation operating. Complete filling and maintain agitation until spraying is complete.

Compatibility: Concero® is compatible with commonly used fungicides and insecticides. Seek advice if mixing with herbicides and/or plant growth regulators.

IT IS AN OFFENCE FOR USERS OF THIS PRODUCT TO CAUSE RESIDUES EXCEEDING THE RELEVANT MRL IN THE FOOD NOTICE: MAXIMUM RESIDUE LEVELS FOR AGRICULTURAL COMPOUNDS.

Withholding Periods:

Barley, wheat and triticale (grain) - 56 days

Wheat, barley, triticale (forage) - 42 days

Resistance Management: Concero® contains bixafen from the SDHI (Succinate DeHydrogenase Inhibitor) mode of action (Group 7) and prothioconazole from the DMI (DeMethylation Inhibitor) mode of action (Group 3). Resistance to this fungicide and related products could develop in some disease situations from repeated use. To minimize this risk use a maximum of 2 applications per season. Make no more than 2 applications using fungicides from the SDHI group and no more than 3 applications, including seed treatment, from the DMI group.

Registered to and distributed by: Adria New Zealand Limited. P.O. Box 535 Kumeu, Auckland 0841, NEW ZEALAND.

Ph: +64-9-412-9817; www.adria.nz. Concero® is a Registered Trademark and is Registered pursuant to the ACVM Act 1997, Approval No. P10429. EPA Approval Code: HSR101566.