

WARNING

KEEP OUT OF REACH OF CHILDREN

BEETUP COMPACT[®]

**Controls weeds in fodder,
sugar and red beets.**

GROUP C1 HERBICIDE

Registered pursuant to the ACVM Act 1997 No.P5493.
See www.foodsafety.govt.nz for registration conditions.

In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in an emergency only,
call 0800 734 607 (24 hours).

**Herbicide
Contains:
80g/litre
phenmedipham
plus 80g/litre
desmedipham
in the form of
an emulsifiable
concentrate.**



9 421903 419422 >

Net Contents : 5 litres



HSNO Classifications: 3.1D, 6.1D, 6.3B, 6.4A, 6.5B, 6.9B, 9.1A, 9.2A, 9.3C Approved under the HSNO Act 1996, HSR100825. See www.epa.govt.nz for controls.

May be harmful if swallowed, inhaled or absorbed through or in contact with the skin. May cause damage to organs from single, prolonged or repeated exposure. Causes mild skin irritation. Causes serious eye irritation. May cause sensitisation from prolonged skin contact. May cause an allergic reaction to skin. Combustible liquid.

PRECAUTIONS: Store safely away from children and foodstuffs. Avoid contamination of seeds, fertilisers and other pesticides. Avoid contact with skin, eyes and inhalation of spray mist. When mixing and applying wear eye protection, impervious gloves, boots and cotton overalls. Wash contaminated clothing before reuse. Avoid contamination of any water supply with the chemical or empty container.

COMBUSTIBLE LIQUID: Keep away from open flames and heat sources. No smoking. Use water spray or fog to extinguish fire. Do not use water jet.

STORAGE: Store in original container, tightly closed and in a locked dry, cool, well ventilated area away from foodstuffs.

ENVIRONMENTAL INFORMATION: Very toxic to the soil environment. Harmful to terrestrial vertebrates. **VERY TOXIC TO AQUATIC LIFE WITH LONGLASTING EFFECTS.**

Avoid contamination of any water supply or source with chemical or empty container.

SPILLS: Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers.

Wash area with water and absorb with further inert material. Dispose of waste safely (such as to a suitable landfill).

Container Disposal: Triple rinse container and add residue to spray tank. Return empty container to an AgRecovery collection point for disposal.



FIRST AID: If swallowed do NOT induce vomiting. For advice, contact the National Poisons Centre on 0800 POISON (0800 764766) or a doctor immediately. If exposed or if you feel unwell: Call a POISON CENTRE or doctor/physician.

Eye contamination: If product or spray enters eyes, wash out immediately and continue rinsing with clean water for at least 15 minutes. Obtain medical attention.

If eye irritation persists: Get medical advice/attention.

Skin contact: After contact with skin, wash immediately with soap and water. If skin irritation or rash occurs: Get medical advice/attention.

Batch number and manufacturing date: See on packaging

Notice to Buyer: Careful tests have proved that the product is suitable for the recommended purposes when used in accordance with our instructions. However, since storage and use of product are beyond our control, we do not therefore accept any liability for damage or loss of efficacy which may result from the use of this product, whether used in accordance with the directions or not. We are liable for a consistent quality of the product, but the risk of its storage and use is not borne by us.

Registered to: UPL NZ Ltd. Level 6, WHK Tower, 51-53 Shortland St, Auckland 1010.

Distributed by: Adria New Zealand Ltd. 407 State Highway 16, Kumeu, Auckland. Ph: 09 412 9817. Fax: 09 412 9807. www.adria-nz

General Information: Beetup Compact herbicide acts through the foliage of seedling broadleaved weeds to give selective weed control in sugar and red beet crops. This product should be used strictly in accordance with label recommendations.

Susceptible Weeds Include: Seedling stages of chickweed, fathen, fumitory, groundsel, redroot, shepherd's purse, spurrey, nettle and twin cress.

Black nightshade, cornbind and willow weed are controlled only if treated up to the four true leaf stage.

Resistance Management: Beetup Compact is a **GROUP C1 HERBICIDE** from the Phenyl Carbamate herbicide group. Resistance to this herbicide and related products could develop in some situations from repeated use. To minimise this risk, use strictly in accordance with the label instructions. For detailed resistance management strategies refer to: <http://resistance.nzpps.org>.

Resistant Weeds Include: Grass weeds, mature annual broadleaf weeds, perennial broadleaf weeds and wireweed.

Directions for Use: Beetup Compact does not require the addition of surfactants. Ensure spray equipment is thoroughly cleaned before use. Do not apply in more than 280 litres of water per hectare.

Mixing: Partly fill the spray tank and add the required amount of Beetup Compact herbicide with agitation system operating. Add the remaining water and any mixing partners.

Maintain agitation until spraying is complete. Do not allow spray mixture to stand overnight. Flush equipment with clean water after use.

Beetup Compact herbicide may be broadcast or band treated. For band treatment ensure that the use rate per sprayed area is consistent with label recommendations.

Recommendations for Use:

CROP	TIMING	RATE	REMARKS
Sugar Beet and Red Beet	After the crop has 2 true leaves fully developed.		Damage may occur especially on Red Beet if treated prior to 2 true leaf stage.
	Weeds up to 4 true leaf.	6 litres/ha in 200 litres of water.	Best results are obtained when weeds are treated at the youngest possible stage.
	Weeds from 4-6 true leaves and hard to kill weeds up to 4 true leaves.	9 litres/ha in 200 litres of water.	A repeat application may be made to late germinating weeds at least 7 days after the first treatment.
Fodder Beet	Apply after the crop has 2 true leaves fully developed.	3 litres/ha in 200-300 litres of water.	Best results are obtained by applying Beetup Compact following a pre-emergence application of Ethofumesate in mixture with Mitron 70WG. Make the first application of Beetup Compact in mixture with Ethofumesate (50% w/v) at 2 litres/ha plus Mitron 70WG at 1.5kg/ha when weeds are from cotyledon to 2 true leaf stage and a second application 10-14 days later.

NOTE: DO NOT MIX BEETUP COMPACT WITH ANY FOLIAR NUTRIENTS OR SURFACTANTS.

IT IS AN OFFENCE FOR USERS OF THIS PRODUCT TO CAUSE RESIDUES EXCEEDING THE RELEVANT MRL IN THE FOOD NOTICE: MAXIMUM RESIDUE LEVELS FOR AGRICULTURAL COMPOUNDS.

Withholding Period: Fodder beet – 56 days. Do not graze stock on treated areas for 56 days after spraying.

Additional Information: Do not apply if air temperature is likely to rise above 29°C, and avoid treatment if rain is likely before spray has dried on the foliage. Herbicide activity will be slowed down in cool weather.