



Concord[®] 300SC

Fungicide for the control of Early and Late Blight in Potatoes and Downy mildew in Onions and Garlic.

Active Ingredient: 300g/litre cymoxanil

Chemical Group: Acetamide

Formulation: Suspension concentrate

Pack Size: 5 Litre

Compatibility: Concord is compatible with most fungicides and insecticides. Not compatible with Bordeaux mixture or lime sulphur or other alkaline products.

Concord[®] 300SC possesses excellent translaminar and penetrant ability in fighting Late blight and Downy mildew. Concord has strong curative activity, with activity up to 2 days after infection. It has multi-site activity, with a very low risk level of resistance development.



Powerful curative activity ✓

No known resistance issues ✓

Strong performance under high-pressure disease conditions ✓

Preventative and anti-sporulant activity ✓

Very cost-effective ✓

General Information: Concord is a contact and systemic fungicide with curative activity against downy mildew and late blight when applied immediately after spore germination.

Resistance Management: Concord belongs to the cyanoacetamide chemical group. Diseases resistant to this group of fungicides may develop if overused. To minimise the development of resistance, use strictly in accordance with label recommendations. Always apply in mixture with a mancozeb fungicide. Do not use more than 3 times in succession and do not apply more than 5 times per season.

Compatibility: Concord is compatible with most fungicides and insecticides. Not compatible with Bordeaux mixture or lime sulphur or other alkaline products.

IT IS AN OFFENCE FOR USERS OF THIS PRODUCT TO CAUSE RESIDUES EXCEEDING THE RELEVANT MRL IN THE NEW ZEALAND (MAXIMUM RESIDUE LIMITS OF AGRICULTURAL COMPOUNDS) FOOD STANDARDS.

Withholding Periods:

Onions, Garlic and Potatoes: 14 days

IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS.

CROP	DISEASE	REMARKS
Onions Garlic	Downy Mildew	Apply 370-550ml/ha in tank-mixture with mancozeb at onion/garlic label rates. Use in a programme with suitable alternative fungicides. Begin spraying at the 2 true leaf stage and continue on a 7-10 day schedule, using Concord during periods of high disease pressure. Do not use more than 3 times in succession and do not apply more than 5 times per season.
Potatoes	Late Blight Early Blight	Use 330-380ml/ha in tank-mixture with mancozeb at potato label rates. Use during periods when disease pressure is high and use alternative fungicides at other times. Do not use more than 3 times in succession and do not apply more than 5 times per season. Use the higher rate under extreme infection conditions (warm with high humidity).