

# WARNING

KEEP OUT OF REACH OF CHILDREN

# Corona<sup>®</sup> 250WG

**A systemic FUNGICIDE for the control of Brownrot in summerfruit.**

Active Ingredient: Contains 250g/kg tebuconazole in the form of a water dispersible granule.

Registered pursuant to the ACVM Act 1997 No. P7813


See [www.nzfsa.govt.nz/acvm/](http://www.nzfsa.govt.nz/acvm/) for registration conditions

HSNO Classifications: 6.9B, 9.1A

**This product must be under the care of an Approved Handler during use.**

In a transport emergency dial 111, Police or Fire Brigade.

For specialist advice in a First Aid emergency only call: National Poisons Centre on 0800 764 766.



**ECOTOXIC  
DANGEROUS TO THE ENVIRONMENT**

## Net Contents: 1 Kg

**READ THIS LABEL COMPLETELY BEFORE USE**

**WARNING:** This substance may be harmful if swallowed, inhaled or absorbed through the skin. May cause internal organ damage from repeated oral exposure at high doses.

**PRECAUTIONS:** Keep out of reach of children. Wear protective clothing such as impervious gloves, waterproof hat, coat and trousers when using. Avoid contact with skin or eyes and inhalation of spray mist. Wash hands and exposed skin after use and before meals. When mixing or apply wear waterproof gloves.

**CONTAINER DISPOSAL:** Triple rinse container and add residue to spray tank. Burn, if circumstances, especially wind direction permit, otherwise bury in landfill.

**VERY TOXIC TO AQUATIC ORGANISMS WITH LONG-LASTING EFFECTS.** Avoid contamination of any water supply with chemical or empty container. A strategy to minimise spray drift should be employed at all times when applying spray near aquatic environments.

**FIRST AID:** If swallowed do NOT induce vomiting. For advice, contact the National Poisons Centre on 0800 POISON (0800 764766) or a doctor immediately.

**Eye contamination:** If product or spray enters eyes, wash out immediately and continue rinsing with clean water for at least 15 minutes. Obtain medical attention.

**Skin contact:** After contact with skin, wash immediately with soap and water.

**SPILLS:** Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely, according to local Council regulations.

**Notice to Buyer:** Careful tests have proved that the product is suitable for the recommended purposes when used in accordance with our instructions. However, since storage and use of product are beyond our control, we do not therefore accept any liability for damage or loss of efficacy which may result from the use of this product, whether used in accordance with the directions or not. We are liable for a consistent quality of the product, but the risk of its storage and use is not borne by us.

**Registered to and Distributed by:** Adria New Zealand Ltd.  
P O Box 535, 407 State Highway 16, Kumeu, Auckland.  
Phone: 09 412 9817 Fax: 09 412 9807



**Batch number and manufacturing date:** See on packaging.

**Storage:** Store in original container tightly closed and in a locked, dry, cool, well ventilated area away from foodstuffs. When stored appropriately this product should show no significant degradation for 2 years from the date of manufacture. Contact your supplier for further information about the use of the product that is older than this.

**DIRECTIONS FOR USE:** Apply as a dilute spray, ensuring complete coverage, without excessive run-off. For concentrate spraying, adjust dilution rate accordingly.

CROP	DISEASE	RATE	CRITICAL COMMENTS
Summerfruit	Brownrot	40g/100 litres of water. Use minimum of 800g/ha.	<b>A:</b> Apply in a regular blossom programme from pink to shuck fall inclusive. Thereafter apply alternative non-DMI fungicides through until harvest. or <b>B:</b> Apply 2-3 sprays at 7-10 day intervals in the period prior to harvest. When using pre-harvest use alternative non-DMI fungicides earlier in the season.

Apply a maximum of 3 applications, preferably with non-DMI fungicides, as required over the pink to shuck fall period. Thereafter apply alternative non-DMI fungicides through until harvest.

**Mixing:** Add the required amount of Corona 250WG into the half-filled spray tank with agitation operating. Continue agitation during spraying and after a stoppage before recommencing spraying.

**Application:** Make sure that spray equipment is correctly calibrated. Ensure complete spray coverage for best results. Do not overspray.

**Compatibility:** Corona 250WG is compatible with most of the commonly used insecticides and fungicides, except alkaline materials such as Bordeaux mixture and lime sulphur.

**Withholding Period:**  
Summerfruit – 1 day

**Resistance Management:** Resistance to Corona 250WG and related DMI fungicides could develop from repeated use. To minimise this risk, use strictly in accordance with label instructions. DO NOT make more than 3 applications during the season.