

DIRECTIONS FOR USE:

**Mixing:** Shake well before use. Add the required amount to a partly filled spray tank with agitation system operating. Complete filling the tank with water and apply. Flush spray system with clean water after use.

**Compatibility:**

DO NOT USE CORONET IN ANY MIXTURES WHEN APPLYING TO PERENNIAL RYEGRASS.

When used on wheat, Coronet can be tankmixed with Accuro 125SC and chlormequat-chloride in 2 and 3 way mixtures. Care must be taken to ensure that recommended timings for all products are followed. The addition of a suitable adjuvant oil at label rates is recommended. Do not use Coronet within 5 days of broad-leaved herbicide applications.

CROP	WEEDS	RATE	REMARKS
Wheat	Wild oats	750mL/ha in 250 litres of water. Add adjuvant oil at recommended rates.	Apply when wild oats are between 3 leaf and 1st node visible growth stage and when they are growing actively.
	Lesser canary grass ( <i>Phalaris minor</i> )	750mL/ha in 250 litres of water. Add adjuvant oil at recommended rates.	Apply when weeds are between the 3 leaf visible stage and the end of tillering and are growing actively. Avoid application during periods of moisture stress.
Perennial ryegrass ( <i>Lolium perenne</i> )	Wild oats	750mL/ha in 250 litres of water. <b>Do not use with any other agrochemicals or adjuvants.</b>	Apply in early spring when wild oats are between 3 leaf and end of tillering growth stage and are actively growing. Note: Seed yield response is likely to be reduced the longer the delay in the removal of wild oats in spring.

DO NOT USE ON BARLEY

IMPORTANT NOTES:

Gnawed canary grass (*Phalaris paradoxa*) is not controlled by Coronet at label rates.

Do not overlap swath width in ryegrass crops as yield suppression may occur.

Do not use Coronet with any other agrochemical or spray additive when applied to Perennial ryegrass.

Resistant weeds: Annual and Hybrid ryegrasses, tall fescue, chewing's fescue, cocksfoot, browntop, grazing brome (=spiky brome – Bromus stamineus), prairie grass and most broadleaved weeds.

Rainfastness: Coronet is rainfast within one hour of application.

**IT IS AN OFFENCE for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**

**Withholding periods:**

Wheat - 70 days (Do not graze treated crop for at least 70 days after treatment)

Perennial ryegrass - 42 days (Do not graze treated crop for at least 42 days after treatment)

**Soil Activity:** Coronet is rapidly broken down by soil microbes causing no restrictions on crop rotations.

Undersown crops: Coronet can be used on crops undersown with certain grasses (See under "Resistant Weeds" above) and clover.

Do not use on crops undersown with "Concord" ryegrass.

WARNING  
KEEP OUT OF REACH OF CHILDREN

Coronet®

A selective grass herbicide for use on autumn and spring sown wheat and perennial ryegrass for the control of wild oats and lesser canary grass (*Phalaris minor*).

Active Ingredient: contains 69 g/litre fenoxaprop-P-ethyl in the form of an oil-in-water emulsion.

GROUP A HERBICIDE

Registered pursuant to the ACVM Act 1997 No. P8196  
See www.foodsafety.govt.nz for registration conditions

Must be under the personal control of an approved handler when the substance is applied in a wide dispersive manner; or used by a commercial contractor.

In a transport emergency dial 111, Police or Fire Brigade.  
For specialist advice in a First Aid emergency only call:  
National Poisons Centre on 0800 764 766.



Net Contents: 10 litres



Approved under the HSNO Act 1996, HSR100440. See www.epa.govt.nz for controls.

HSNO Classifications: 6.3B, 6.5B, 6.9B, 9.1A

Records should be kept of the use and disposal of this product.

This substance may be harmful if swallowed, in-haled or absorbed through the skin, and may cause irritation to eyes and skin. May cause an allergic skin reaction. Avoid inhalation of vapours or mist from spray. May cause internal organ damage from single or repeated oral exposure.

**Precautions:** Store in original container, tightly closed, away from foodstuffs in a cool dry, secured area out of direct sunlight. Avoid contaact with skin and eyes, and inhalation of spray mist. Do not eat, drink or smoke while using this product. Wear protective gloves, protective clothing, eye protection and face protection.Wash hands and exposed skin before meals and after work. Contaminated clothing should not leave the workplace and must be cleaned before re-use. Avoid contaminati-on of any water supply with chemical or empty container. Apply this product carefully. Spray drift may cause serious damage to desirable plants. Ensure compliance with relevant regional air management plans.

**VERY TOXIC TO AQUATIC ORGANISMS WITH LONGLASTING EFFECTS.** A strategy to minimise spray drift should be employed at all times when applying spray near aquatic environments.

**Disposal of Container:** Triple rinse container and add residue to spray tank. Return clean, empty container to an AgRecovery site for disposal. Avoid contamination of any water supply with chemical or empty container.



**FIRST AID: Ingestion:** Do NOT induce vomiting if swallowed. Wash mouth with water and seek medical attention immediately. Small sips of water may be given. For advice, contact the National Poisons Centre on 0800 POISON (0800 764766). **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Seek medical attention immediately. **Skin Contact:** Immediately flush body and clothes with large amounts of water. Remove contaminated clothing and footwear. Gently wash affected areas with soap and water. If a large area is affected or if irritation or rash persists seek medical assistance. **Inhalation:** Should irritation occur, remove patient to fresh air. If breathing is shallow or has stopped, ensure airway is clear and apply resuscitation. Seek medical assistance soon if irritation persists.

**SPILLS:** Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely (such as to a suitable landfill).

**Batch No. and Manufacturing Date:** See on packaging

**Registered to and Distributed by:** Adria New Zealand Limited.  
P.O. Box 535, 407 State Highway 16 Kumeu, Auckland.  
Ph: 09 412 9817. Fax: 09 412 9807. www.adria.nz.



**Notice to Buyer:** Careful tests have proved that the product is suitable for the recommended purposes when used in accordance with our instructions. However, since storage and use of product are beyond our control, we do not therefore accept any liability for damage or loss of efficacy which may result from the use of this product, whether used in accordance with the directions or not. We are liable for a consistent quality of the product, but the risk of its storage and use is not borne by us.

**Resistant Weeds Warning:** Coronet is a **GROUP A HERBICIDE** from the aryloxyphenoxypropionates (FOPs) chemical group. Naturally occurring weed biotypes resistant to aryloxyphenoxypropionate herbicides can become more prevalent after many years of continuous use. To delay the onset of, or control resistant weeds, use in rotation with herbicides of a different mode of action. Since the occurrence of resistant weeds is difficult to detect prior to use, Adria New Zealand Limited accepts no liability for losses that may result from the failure of this product to control resistant weeds.