

Curator®

For the control of certain diseases in mushrooms, ornamentals and stonefruit.



Active Ingredient: 462g/kg prochloraz as the manganese complex

Chemical Group: Imidazole

Formulation: Wettable Powder

Pack Size: 1 kg

Curator offers highly cost-effective disease control, with many key benefits:

- ✓ Excellent activity against *Verticillium* spp. In mushrooms.
- ✓ Broad spectrum disease control in ornamentals with a high degree of crop safety.
- ✓ Good tank-mixing flexibility .
- ✓ Reliable performance and highly cost-effective disease control.

About Curator®:

Curator is a high-quality powder formulation of prochloraz.

Curator is a proven performer in mushrooms, ornamentals and stonefruit, and controls a wide spectrum of diseases. Curator is non-systemic but has strong contact activity and translaminar mobility.

General Information:

Curator is a DMI fungicide which may be applied as a spray treatment to ornamentals and stonefruit, as a dip treatment to ornamentals, and a drench treatment in mushrooms.

Resistance Management: Resistance to this and other DMI products could develop from repeated use. To minimise this risk use strictly in accordance with label instructions.

Directions for use:

The method of application shall be limited to ground based application only:

Mixing: Thoroughly pre-mix the required amount with a little water to form a slurry. Add water whilst agitating. For spray treatments maintain agitation while mixing and applying. Also agitate thoroughly after a stoppage, before recommencing spraying. Use immediately after mixing.

Compatibility: Curator is compatible with most commonly used insecticides and fungicides. DO NOT use with alkaline materials such as lime sulphur or Bordeaux mixture.

Recommendations: Rates suggested are for high volume spraying to run-off. For concentrate spraying, adjust dilution rate accordingly.

Crop	Disease	Rate	Critical Comments
Mushrooms	Brown spot (Verticillium spp.)	3g/square metre in 1 litre of water as a drench treatment.	Apply 7-9 days after casing. Do not apply within 10 days of picking. Apply a maximum of 3 applications per season.
Ornamentals	Colletotrichum spp., Glomerella leaf spot, Monochaetia spp., Pestalotia spp., Powdery mildew	100g/100 litres in sufficient water to obtain thorough spray coverage.	Apply at first signs of infection and repeat at 14 day intervals.
		100g/100 litres as a dip.	Immerse whole plants in dip treatment prior to shipping. Apply a maximum of 3 applications per season
Stonefruit	Brown rot	25g/100 litres	Apply at pink and repeat at 7-10 day intervals until petal fall. Shorter intervals may be required in wet weather. Use in mixture with non-DMI fungicides and apply no more than three treatments per season. Apply a maximum of 3 applications per season.

IT IS AN OFFENCE FOR USERS OF THIS PRODUCT TO CAUSE RESIDUES EXCEEDING THE RELEVANT MRL IN THE FOOD NOTICE (MAXIMUM RESIDUE LEVELS FOR AGRICULTURAL COMPOUNDS).

Withholding Periods:

Mushrooms: 10 days

Stonefruit: Do not use after petal fall.

Registered to and distributed by: Adria New Zealand Limited. P.O. Box 535 Kumeu, Auckland 0841, NEW ZEALAND.
Ph: +64-9-412-9817 www.adria.nz.

Curator is Registered pursuant to the ACVM Act 1997, Approval No. P7738. Approved under the HSNO Act 1996, HSR007983.

For more information call us on +64-9-412-9817 | www.adria.nz