



# Curator

For the control of certain diseases in mushrooms, ornamentals and stonefruit.

Active Ingredient: 462g/kg prochloraz as the manganese complex.

Chemical Group: Imidazole

Formulation: Wettable Powder

Pack Size: 1 kg

Compatibility: Curator is compatible with most commonly used fungicides and insecticides. DO NOT use with alkaline materials such as lime sulphur or Bordeaux mixture.

Curator is a new high-quality formulation of 'prochloraz', and is the first generic competitor to Octave®<sup>1</sup> to be registered in NZ.

Curator is a proven performer in mushrooms, ornamentals and stonefruit, and controls a wide spectrum of diseases. Curator is non-systemic but has strong contact activity and translaminar mobility.



Curator offers highly cost-effective disease control, with many key benefits:

- Excellent **activity** against **Verticillium spp.** In **mushrooms** ✓
- **Broad spectrum** disease control in **ornamentals** with a high degree of **crop safety** ✓
- **Good tank-mixing flexibility** ✓
- Reliable performance and highly **cost-effective** disease control ✓

**GENERAL INFORMATION:**

Curator is a DMI fungicide which may be applied as a spray treatment to ornamentals and stonefruit, as a dip treatment to ornamentals, and a drench treatment in mushrooms.

**RESISTANCE MANAGEMENT:** Resistance to this and other DMI products could develop from repeated use. To minimise this risk use strictly in accordance with label instructions.

**DIRECTIONS FOR USE:**

**The method of application shall be limited to ground based application only.**

**Mixing:** Thoroughly pre-mix the required amount with a little water to form a slurry. Add water whilst agitating. For spray treatments maintain agitation while mixing and applying. Also agitate thoroughly after a stoppage, before recommencing spraying. Use immediately after mixing.

**COMPATIBILITY:** Curator is compatible with most commonly used insecticides and fungicides. DO NOT use with alkaline materials such as lime sulphur or Bordeaux mixture.

**RECOMMENDATIONS:** Rates suggested are for high volume spraying to run-off. For concentrate spraying, adjust dilution rate accordingly.

CROP	DISEASE	RATE	REMARKS
Mushrooms	Brown spot ( <i>Verticillium spp.</i> )	3g/square metre in 1 litre of water as a drench treatment.	Apply 7-9 days after casing. Do not apply within 10 days of picking. Apply a maximum of 3 applications per season.
Ornamentals	<i>Colletotrichum spp.</i> , Glomerella leaf spot, <i>Monochaetia spp.</i> , <i>Pestalotia spp.</i> , Powdery mildew	100g/100 litres in sufficient water to obtain thorough spray coverage.  100g/100 litres as a dip.	Apply at first signs of infection and repeat at 14 day intervals.  Immerse whole plants in dip treatment prior to shipping. Apply a maximum of 3 applications per season.
Stonefruit	Brown rot	25g/100 litres	Apply at pink and repeat at 7-10 day intervals until petal fall. Shorter intervals may be required in wet weather. Use in mixture with non-DMI fungicides and apply no more than three treatments per season. Apply a maximum of 3 applications per season.

**IT IS AN OFFENCE FOR USERS OF THIS PRODUCT TO CAUSE RESIDUES EXCEEDING THE RELEVANT MRL IN THE NEW ZEALAND (MAXIMUM RESIDUE LIMITS OF AGRICULTURAL COMPOUNDS) FOOD STANDARDS.**

**WITHHOLDING PERIODS:**

Mushrooms: 10 days

Stonefruit: Do not use after petal fall

Registered to and Distributed by: Adria New Zealand Limited. P.O. Box 535 Kumeu, Auckland 1250, NEW ZEALAND.

Ph: +64-9-412-9817; Fax: +64-9-412-9807

[www.adriacp.co.nz](http://www.adriacp.co.nz)

Curator is Registered pursuant to the ACVM Act 1997, Approval No. P7738

®1Octave is a Registered Trademark of Bayer AG and is registered pursuant to the ACVM Act 1997, Approval No. P3055