

IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS.

DIRECTIONS FOR USE:

CROP	DISEASE	RATE	CRITICAL COMMENTS
Grapes	Powdery Mildew (Uncinula necator)	7.5mL/100L	Apply non-DMI fungicides (eg sulphur) prior to flowering. From flowering apply Cypro 200SC at 2 to 3 week intervals. Ensure complete coverage using not less than 1000 litres water per hectare. Early season Cypro 200SC treatments may cause leaf discolouration on some grape varieties but this has not affected yield or fruit quality.
Peas	Powdery Mildew (Erysiphe pisi)	125mL/ha	Apply as a preventative spray when conditions favour disease development or at the first appearance of disease symptoms. If conditions persist for infection a repeat application may be necessary after 14 days.
Stonefruit	Brown Rot (blossom blight) (Monilinia spp)	7.5mL/100L	Make the first application at pink and repeat at 7 to 10 day intervals until shuck fall. Shorter intervals may be required in wet weather.
Vegetable and Field Brassicas	Ring Spot (Mycosphaerella brassicicola)	200mL/ha plus 2.5 L/ha 'all purpose' emulsifiable crop mineral oil	Apply as a preventative spray when conditions first favour disease development, or at the first sign of disease symptoms. Repeat at 2 to 3 week intervals. Apply a maximum of 4 applications per crop.
Onions, Garlic	White rot	500-750 mL/ha in 1000 litres of water/ha	Make the first application 6-8 weeks after sowing. Make 2 further applications at 4 week intervals.
	Rolf's disease		Apply in early December as either the third White rot application or as a fourth application depending on timing of previous sprays.

TURF GRASS DISEASE CONTROL

DISEASE	RATE PER 100M ²	APPLICATION INTERVAL	CRITICAL COMMENTS
Brown Patch (Rhizoctonia solani)	15 – 20ml	21–28 days	Apply preventatively when conditions favour disease development, especially when warm temperatures and high humidity persist. When disease pressure is high use the shorter application interval and/or the higher rate. Repeat as necessary. DO NOT apply more than 4 applications of Cypro 200SC per year. Integrate with good turf management and a spray program with other fungicides. DO NOT apply more than 15 mL/100 square metres to creeping bent (Agrostis stolonifera) or Poa annua at intervals shorter than 28 days as phytotoxicity may occur. DO NOT apply to Cotula greens. Apply with properly calibrated equipment in 1000 to 2000 litres water/ha.
Dollar Spot (Sclerotinia homeocarpa)	10ml	21–28 days	DO NOT apply more than 15 mL/100 square metres to creeping bent (Agrostis stolonifera) or Poa annua at intervals shorter than 28 days as phytotoxicity may occur. DO NOT apply to Cotula greens. Apply with properly calibrated equipment in 1000 to 2000 litres water/ha.
Rusts (Puccinia spp)	10ml	28 days	DO NOT apply more than 15 mL/100 square metres to creeping bent (Agrostis stolonifera) or Poa annua at intervals shorter than 28 days as phytotoxicity may occur. DO NOT apply to Cotula greens. Apply with properly calibrated equipment in 1000 to 2000 litres water/ha.
Red Thread (Laetisaria fuciformis)	15 – 20ml	21–28 days	DO NOT graze animals on treated turf areas, or use grass clippings for feed.

MIXING: Add the required amount of Cypro 200SC to the half-filled spray tank. Add the remainder of the water and agitate thoroughly while mixing, spraying and when restarting after a stoppage.

COMPATIBILITY: Cypro 200SC is compatible with sulphur, mancozeb, streptomycin and most other fungicides and insecticides. Ensure thorough agitation of the mixture while spraying. DO NOT tank mix Cypro 200SC with copper fungicides.

IT IS AN OFFENCE FOR USERS OF THIS PRODUCT TO CAUSE RESIDUES EXCEEDING THE RELEVANT MRL IN THE FOOD NOTICE: MAXIMUM RESIDUE LEVELS FOR AGRICULTURAL COMPOUNDS.

Withholding Periods: Grapes – 28 days; Peas – 14 days; Stonefruit – DO NOT apply after shuckfall; Brassicas – 14 days; Onions, Garlic – 21 days.

UN 3082. Class: 9, Packaging group: III. ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (CYPROCONAZOLE SOLUTION).

Do not carry this product on a passenger service vehicle.

WARNING

KEEP OUT OF REACH OF CHILDREN

Cypro[®] 200SC

**For the control of diseases in turf,
fruit and vegetable crops.**

Active Ingredient: Contains 200g/litre cyproconazole
in the form of a suspension concentrate.

GROUP 3 FUNGICIDE

Registered pursuant to the ACVM Act 1997 No. P9401
See www.foodsafety.govt.nz for registration conditions.

In a transport emergency dial 111, Police or Fire Brigade.
**For specialist advice in an emergency only call:
0800 734 607 (24 hours).**



Net Contents: 1 litre



READ THIS LABEL COMPLETELY BEFORE USE



HSNO Classifications: 6.1D (O), 6.3B, 6.8B, 6.9B, 9.1A, 9.3C.
Approved under the HSNO Act 1996, HSR101190.
See www.epa.govt.nz for controls.
Qualified/Competent Person: Required.

This material is harmful if swallowed. Suspected of damaging fertility or the unborn child. May cause internal organ damage from single or repeated exposure. Suspected teratogen: Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Use personal protective equipment as required. Causes mild skin irritation.

ENVIRONMENTAL INFORMATION: VERY TOXIC TO AQUATIC ORGANISMS WITH LONG-LASTING EFFECTS. Avoid contamination of any water supply with chemical or empty container. A strategy to minimise spray drift should be employed at all times when applying spray near aquatic environments. Harmful to terrestrial vertebrates.

PRECAUTIONS: Store locked up in original container, tightly closed, away from foodstuffs. When handling, mixing or applying wear protective clothing, including face shield, chemically resistant gloves and footwear. Avoid contact with skin and eyes and inhalation of the spray mist. Do not breathe fumes, mist or spray. Wash splashes of concentrate from the eyes or skin immediately. If clothing becomes contaminated with product, remove clothing immediately. DO NOT eat, drink or smoke while using. Wash hands and face before eating, drinking or smoking and after work.

STORAGE AND USE: Store locked up in the original, unopened container in a cool, dry place, out of direct sunlight and away from stockfeed or foodstuffs. As a Class 9 Substance with Ecotoxicity Classifications, storage must be carried out in such a manner as to prevent contamination of waterways. It is recommended that The New Zealand Standard for the Management of Agrichemicals (NZS8409) is followed. See Safety Data Sheet for further information. Apply with well maintained and calibrated spray equipment. Records of use must be kept. Application, storage and record keeping should be as described in the Management of Agrichemicals NZS8409. DO NOT carry on a passenger service vehicle.

APPLICATION CONTROLS:

- 1) This product must not be applied into or onto water.
- 2) This product must be applied by ground based methods only.
- 3) This product must not be applied at rates exceeding 2L/ha per application (400g/ha cyproconazole).
- 4) This product must not be applied to the same area more than 4 times per year.

CONTAINER DISPOSAL: Triple rinse container and add residue to spray tank. Return empty container to an AgRecovery collection point for disposal.



FIRST AID: If Swallowed: Do NOT induce vomiting. Rinse mouth. For advice, contact the National Poisons Centre on 0800 POISON (0800 764 766) or a doctor if you feel unwell. If in eyes: Flush eyes immediately and continue for at least 15 minutes. Seek medical attention. If on skin: Immediately flush body and clothes with large amounts of water. Wash affected areas with soap and water. If skin irritation occurs, get medical advice/attention. If inhaled: Remove patient to fresh air. If breathing is shallow or has stopped ensure airway is clear and apply resuscitation. Seek medical assistance soon if irritation persists. If medical advice is needed, have product container or label at hand. Get medical advice/attention if you feel unwell.

SPILLS: Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely, according to local Council regulations.

SAFETY DATA SHEET: If additional hazard or safety information is required, refer to the Safety Data Sheet. For a copy visit our website at www.adria.nz.

Notice to Buyer: Careful tests have proved that the product is suitable for the recommended purposes when used in accordance with our instructions. However, since storage and use of product are beyond our control, we do not therefore accept any liability for damage or loss of efficacy which may result from the use of this product, whether used in accordance with the directions or not. We are liable for a consistent quality of the product, but the risk of its storage and use is not borne by us.

Registered to and Distributed by:

Adria New Zealand Limited, P.O. Box 535, 407 State Highway 16 Kumeu, Auckland.
Ph: 09 412 9817. Fax: 09 412 9807. www.adria.nz

Batch number and manufacturing date: See on packaging

Resistance Management: Cypro 200SC is a **GROUP 3 FUNGICIDE** from the DeMethylation (DMI/Triazole) group of fungicides. Resistance to Cypro 200SC and related DMI fungicides could develop from repeated use. To minimise this risk, use strictly in accordance with label instructions. **Grapes:** Use a maximum of 4 sprays, preferably in mixture with an approved protectant fungicide. No more than 2 DMI fungicides should be applied alone per season. **Stonefruit:** Use a maximum of 3 applications preferably in mixture with a non-DMI fungicide. Restrict use of the DMI fungicides to one part of the growing season, either during flowering or pre-harvest. Ensure full coverage of foliage.

