



# Cyrus<sup>®</sup>

A systemic INSECTICIDE for the control of thrips in onions and cabbage aphids in vegetable brassicas and aphids in lettuce (cell transplants).

---

<u>Active Ingredient:</u>	350g/litre imidacloprid
<u>Chemical Group:</u>	Chloronicotinyl
<u>Formulation:</u>	Suspension concentrate
<u>Pack Size:</u>	1 and 5 litre

---

Cyrus<sup>®</sup> contains the active ingredient '*imidacloprid*' - a systemic protectant insecticide which is highly active against thrips and aphids. It is generally active on thrip and aphid populations which exhibit resistance to many alternative chemical groups. It acts primarily as a stomach and contact insecticide.



## **Cyrus<sup>®</sup> offers highly cost-effective thrip and aphid control:**

- Systemic activity ✓
- Unique mode of action ✓
- Good IPM fit ✓

**Cyrus** is a systemic insecticide of the chloro-nicotinyl group for the control of thrips in onions, cabbage aphids in vegetable brassicas and aphids in lettuce (cell transplants).

## **DIRECTIONS FOR USE:**

**Mixing:** Pour **Cyrus** directly into the spray tank with agitators running.

### **Resistance Management:**

Resistance to this insecticide could develop in some situations from repeated use. Apply a maximum of four consecutive applications per season according to the Onion thrips management strategy issued by the N Z Onion Exporters Association.

## **IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS.**

### **Recommendations:**

<b>Crop</b>	<b>Pest</b>	<b>Rate</b>	<b>Remarks</b>
Onions	Thrips	225ml/ha	Apply first application when monitoring of thrip levels indicates the threshold level has been reached. Repeat applications at 7-10 day intervals as a cluster of a maximum of 4 applications. Use in 500-600 litres of water/ha. The addition of a wetting agent is suggested. Ensure good spray coverage. Refer to the Resistance Management statement.
Vegetable Brassicas (cell transplants)	Cabbage aphids	9ml/2.5 litres of water per 1000 plants.	Apply evenly in a watering can to seedling trays 1 day before transplanting.
Lettuce (cell transplants)	Aphids	20-30ml/2.5 litres of water per 1000 plants.	Apply evenly in a watering can to seedling trays 1 day before transplanting.

**Compatibility:** **Cyrus** is compatible with commonly used fungicides.

**IT IS AN OFFENCE FOR USERS OF THIS PRODUCT TO CAUSE RESIDUES EXCEEDING THE RELEVANT MRL IN THE NEW ZEALAND (MAXIMUM RESIDUE LIMITS OF AGRICULTURAL COMPOUNDS) FOOD STANDARDS.**

### **Withholding Periods:**

Onions - 7 Days

Vegetable Brassicas (cell transplants) - 70 Days

Lettuce (cell transplants) - 35 Days