

It is an offence to use this product on animals. Unless otherwise specified rates are suggested for high volume spraying to just before run-off. For concentrate spraying, adjust dilution rate accordingly.

Crop	Pest	Rate	Remarks
Beans	Looper caterpillar White butterfly caterpillar	400ml/ha in 500-600 litres of water or 40ml/100 litres.	Apply at first sign of damage and repeat as necessary.
Brassicas	Diamond back moth White butterfly caterpillar	400ml/ha in 700 to 1000 litres of water or 40ml/100 litres.	Apply at 10-14 day intervals from transplanting. The addition of a suitable spreader-sticker is recommended. Refer to Resistance Management comments.
Eucalyptus spp	Tortoise beetle	Combine 500ml with 500ml Butyl Dioxitol. Apply 1 litre of mixture/ha.	Apply by aerial spraying using micronair atomisers.
Tree seedling nursery beds (Pinus spp)	Thrips, Looper caterpillar Tomato fruitworm	40ml/100 litres	Apply every 2-3 weeks from seedling emergence. The addition of a spraying oil at 1% is recommended.
Pre-flowering only. Kiwifruit, Tamarillos and grape vines. Non-bearing pipfruit, stonefruit and avocados. Newly planted shelter provided not in flower.	Grass grub beetle	100ml/100 litres in moderately fine spray.	Ensure complete foliage coverage. Apply at dusk commencing when beetles first appear, and repeat at 6-8 day intervals over the beetle feeding period.
Lucerne	Sitona weevil Aphids	250ml/ha in a minimum of 250 litres of water. 250ml/ha in a minimum of 250 litres of water.	Application should be timed to control adult weevils from mid-May onwards following post-aestivatory flights but prior to egg laying. Apply immediately after cutting if aphids are present. Repeat if necessary.
Onions	Onion thrips	400ml/ha in 500-600 litres of water.	Apply at the first sign of thrip damage and repeat as necessary after 2-3 weeks. The addition of a suitable spreader sticker is recommended. Refer to Resistance Management comments.
Ornamentals	Looper caterpillar Tomato fruitworm, Thrips	40ml/100 litres	Apply at first appearance and repeat as necessary.
Potatoes	Potato tuber moth	500ml/ha in 500-600 litres of water.	To control foliage mining apply at 10-14 day intervals from the first appearance of mining activity. Maintain adequate soil moisture and soil cover over developing tubers to assist in the control of tuber mining.
Tamarillos	White fly	Pre-flowering 80ml/100 litres Post-flowering 50ml/100 litres or 25ml plus 300ml Caltex D-C-Tron NR/100 litres.	Apply immediately after pruning and repeat 7-10 days later. Apply at 10-14 day intervals or as required. Ensure complete coverage of plants, particularly the underside of leaves where whitefly congregate.
Tomatoes (outdoor)	Tomato fruitworm Looper caterpillar	30-40ml/100 litres of water. Apply 400-1000 litres/ha depending on crop density.	Use the lower rate on a 14 day schedule or the higher rate on a 21 day schedule. Begin treatment after emergence or transplanting. For tomato fruitworm control apply only after 7 January and if crop monitoring shows need.
Maize and sweetcorn	Greasy cutworm Corn earworm	500ml/ha 400ml/ha	Apply in evening as a directed spray to thoroughly cover the base of plants. Rain within a few days of treatment will improve control. Apply by air in a minimum of 100 litres of water/ha. Begin at first sign of damage and repeat 14-21 days later or as required.
Winter squash	Aphids	400ml/ha in 500-700 litres of water	Make two applications 10-14 days apart over the pre-flower period. Use alternative insecticides if required for the rest of the season.

For hydraulic knapsack application use 4ml/10 litres. For airblast knapsack sprayers 12ml/10 litres.

UN1993; CLASS 3, PG III, FLAMMABLE LIQUID, N.O.S. (HYDROCARBON SOLVENTS (AROMATIC) (DELTAMETHRIN)). Do not carry this product on a passenger service vehicle.

WARNING
KEEP OUT OF REACH OF CHILDREN

DeltaMax®

A synthetic pyrethroid INSECTICIDE for the control of caterpillars, beetles, thrips and aphids on a broad range of fruit and vegetable crops, and on trees and ornamentals.

Active Ingredient: 25g/litre deltamethrin in the form of an emulsifiable concentrate. Also contains 795g/litre hydrocarbon liquids.

GROUP 3 INSECTICIDE

Registered pursuant to the ACVM Act 1997 No. P8576
See www.foodsafety.govt.nz for registration conditions.

In a transport emergency dial 111, Police or Fire Brigade.
**For specialist advice in an emergency only call:
0800 734 607 (24 hours).**



Net Contents: 5 litre



READ LABEL BEFORE USE



HSNO Classifications: 3.1C, 6.1D, 6.3B, 6.4A, 6.9B, 9.1A, 9.3B, 9.4A. **Approved under the HSNO Act 1996, HSR000328.** See www.epa.govt.nz for controls. **Qualified Person/Contractor:** Required.

WARNING: Flammable liquid and vapour. Harmful if swallowed. Causes mild skin irritation. Causes serious eye irritation. May cause damage to organs through prolonged or repeated exposure. Very toxic to aquatic life with long lasting effects. Toxic to terrestrial vertebrates. Very toxic to terrestrial invertebrates.

SAFETY DIRECTIONS: Keep out of reach of children. Read label before use. Keep away from heat, sparks, open flames or hot surfaces. No smoking. Keep container tightly closed. Take precautionary measures against static discharge. Do not breathe fumes, vapours or spray. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Avoid release to the environment. Wear protective clothing. Collect Spillage. In case of fire: Use dry chemical, water, carbon dioxide or foam for extinction. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area. Do not apply directly into or onto water. Use with adequate ventilation. **TOXIC TO BEES.** Spray must not contact plants in flower if they are likely to be visited by bees. **HANDLING PRECAUTIONS: EQUIPMENT:** Apply using calibrated and maintained equipment in accordance with the NZ Standard for the Management of Agrichemicals (NZS8409).

STORAGE: Store locked up in the original, unopened container in a cool, dry, well-ventilated place, out of direct sunlight and away from stockfeed or foodstuffs. As a Class 9 Substance with Ecotoxicity Classifications, storage must be carried out in such a manner as to prevent contamination of waterways. It is recommended that The New Zealand Standard for the Management of Agrichemicals (NZS8409) is followed. See Safety Data Sheet for further information. **VERY TOXIC TO AQUATIC ORGANISMS.** Avoid contamination of any water supply with product or empty container.

CONTAINER DISPOSAL: Triple rinse container and add residue to spray tank. Return empty container to an AgRecovery collection point for disposal.



FIRST AID: If medical advice is needed, have product container or label at hand. Get medical advice/attention if you feel unwell. **IF SWALLOWED:** Call a POISON CENTER or doctor/physician if you feel unwell. Rinse mouth. **IF ON SKIN (or hair):** Remove/Take off immediately all contaminated clothing. Rinse skin with water/shower. If skin irritation occurs: Get medical advice/ attention. **IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/attention.

SPILLS: Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely (such as to a suitable landfill).

General: DeltaMax is a quick acting synthetic pyrethroid insecticide which works through contact activity and ingestion. It is not systemic. Thorough plant coverage is essential.

Resistance Management: DeltaMax is a **GROUP 3 INSECTICIDE** of the pyrethroid group of chemistry. To minimise the risk of resistance use strictly in accordance with label instructions and resistance management strategies that exist for any insect pest listed on the label. Avoid using this insecticide exclusively all season and avoid unnecessary spraying. Maintain good cultural control practices. For further information, refer to the NZPPS website: <http://resistance.nzpps.org/>

Brassicas – To minimise the development of resistance to synthetic pyrethroids apply only in the period from February to August. Use insecticides of a different mode of action during September to the end of January.

Onions – Refer to supplier or adviser for the local integrated Pest Management programme. A cluster of 3-4 sprays should be made from early January in response to crop monitoring when the threshold of thrip numbers has been exceeded. Ensure good cultural practices such as removal of self-seeds.

Compatibility: DeltaMax is compatible with most commonly used fungicides and insecticides. Flocculation may occur when mixed with some wettable sulphur formulations.

Mixing: Add the required amount of DeltaMax directly into the spray tank with agitator operating. Maintain agitation during application.

IT IS AN OFFENCE for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIODS: Beans, brassicas, tomatoes (outdoor) - 3 days; Onions, potatoes - 14 days

Tamarillos - 7 days; Lucerne - 21 days; Pipfruit, stonefruit, avocados - Non-bearing plants only; Grapes - Use only before the start of flowering; Maize, sweetcorn - 7 days. Do not graze or feed stubble to animals within 14 days of application; Kiwifruit - Do not apply to kiwifruit after bud cracking (first visible petals). Winter squash – Do not use after the start of flowering.

Batch No. and Manufacturing Date: See on packaging.

Registered to and Distributed by: Adria New Zealand Limited.
P.O. Box 535, 407 State Highway 16 Kumeu, Auckland. Ph: 09 412 9817. www.adria.nz



Notice to Buyer: Careful tests have proved that the product is suitable for the recommended purposes when used in accordance with our instructions. However, since storage and use of product are beyond our control, we do not therefore accept any liability for damage or loss of efficacy which may result from the use of this product, whether used in accordance with the directions or not. We are liable for a consistent quality of the product, but the risk of its storage and use is not borne by us.