

DIRECTIONS FOR USE: It is an offence to use this product on animals.

APPLICATION RESTRICTIONS: DO NOT apply more than 1.05 L product per hectare. DO NOT apply more than once a year. Eromite must be applied by ground-based methods only.

SPECIAL NOTES: EroMite is a growth regulator which leads to adult mites laying sterile eggs and prevents existing eggs and nymphs from developing. These stages of the lifecycle are generally controlled within one to two weeks. EroMite does possess some translaminar activity as well as residual control. Subsequently, mite control can last for up to four or five weeks. It should be noted that EroMite will not control adult mites, and these can survive for two to three weeks.

For best results and long term control, EroMite must be sprayed at the first sign of mite crawlers. If there are large numbers of adults, then it will probably be necessary to spray an alternative miticide from a different chemical group to knockdown the adults and lower the chance of any crop damage.

Crop	Pest	Use Rate	Recommendations
Avocados	Six-Spotted mite (SSM)	35 mL/100L	Thorough crop coverage is critical for the best results. Per hectare water rates should not be under 750 litres of water. Concentrate spraying is not advised. For larger trees, per hectare water rates of up to 3,000 litres may be necessary. Note: It is an offence to spray more than 1.05 L/ha of EroMite.
Apples, Pears	European Red mite (ERM), Two-Spotted mite (TSM)		Generally, if more than 15% of the leaves are infested with mites (and there are two or more adult mites per leaf), then an alternative knockdown miticide should be considered. This is very important in pears, where even low mite pressure can be a causal factor in leaf russetting. If further control is necessary, use a registered miticide from an alternative chemical group.

IT IS AN OFFENCE for users of this product to cause residues exceeding the relevant MRL in the Food Notice:

Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIOD: Avocados, Pipfruit: Do not harvest for 14 days after application.

For export markets (except Australia) contact your exporter for more details.

This product shall be applied via ground-based application methods only.

EroMite may be harmful to non-target arthropods, but is deemed safe to bees.

GENERAL RECOMMENDATIONS:

CROP MONITORING: Regular crop monitoring of crops during the season at 3-5 day intervals is essential for achieving effective mite control.

MIXING/APPLICATION: Shake container thoroughly before use.

Apply by ground-based application methods only.

Measure the required volume of EroMite, and add it to the partially filled spray tank, and then add the remainder of the water.

- Use a spray unit which is designed to apply high water volumes up to the point of spray run-off and suited to the crop being sprayed.
- Ensure that the sprayer can achieve consistent coverage throughout the crop canopy. Apply enough water to cover the crop to the point of run-off. Avoid excessive spray run-off.
- The required water rate may be established by applying varying test volumes, and adjusting settings on the sprayer, utilising advice based on industry guidelines and/or expert advice.
- Add the required amount of EroMite recommended in the Directions for Use table for per 100L of water. Spray to the point of run-off.
- As the crop grows, the necessary dilute spray volume will change and the sprayer set up and operation may also need to be adjusted.



UN 3082; CLASS 9, PGIII, ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (ETOXAZOLE SOLUTION). Do not carry this product on a passenger service vehicle.

DANGER
KEEP OUT OF REACH OF CHILDREN

EroMite®

A miticide for controlling European Red mite and Two-Spotted mite in pipfruit and Six-Spotted mite in avocados.

Active Ingredient: Contains 110g/litre etoxazole in the form of a suspension concentrate.

GROUP 10 INSECTICIDE

Registered pursuant to the ACVM Act 1997 No. P9431

See www.foodsafety.govt.nz for registration conditions.

In a transport emergency dial 111, Police or Fire Brigade.

**For specialist advice in an emergency only call:
0800 734 607 (24 hours).**



Net Contents: 5 litres



READ LABEL COMPLETELY BEFORE USE



EPA Approval Code: HSR101214. See www.epa.govt.nz for controls.
Hazard Classifications: 6.3B, 6.9A, 9.1A
Qualified Person/Contractor: Required.

Causes mild skin irritation. Causes damage to organs through prolonged or repeated exposure. Very toxic to aquatic life.

SAFETY DIRECTIONS: Keep out of reach of children. Read label before use. Do not breathe fumes, vapours and spray. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Avoid release to the environment. Do not apply directly into or onto water. Take all reasonable steps to ensure that the substance does not cause any significant adverse effects to the environment beyond the application area. Wear protective clothing. Collect Spillage.

STORAGE: Store locked up in the original, unopened container in a cool, dry place, out of direct sunlight and away from stockfeed or foodstuffs. As a Class 9 Substance with Ecotoxicity Classifications, storage must be carried out in such a manner as to prevent contamination of waterways. It is recommended that The New Zealand Standard for the Management of Agrichemicals (NZS8409) is followed. See Safety Data Sheet for further information.

ECOTOXICITY: This substance may be harmful to non-target arthropods.

APPLICATION: Eromite must not be applied into or onto water.

Avoid contamination of any water supply with product or empty container. A strategy to minimise spray drift should be employed at all times when applying spray near aquatic environments.

DISPOSAL OF CONTAINER: Triple rinse container and add residue to spray tank.

Return clean, empty container to an AgRecovery site for disposal.



FIRST AID: If medical advice is needed, have product container or label at hand. Get medical advice/attention if you feel unwell. If skin irritation occurs: Get medical advice/attention.

SPILLS: Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely (such as to a suitable landfill).

Batch No. and Manufacturing Date: See on packaging

Registered to and Distributed by: Adria New Zealand Limited.

P.O. Box 535, 407 State Highway 16 Kumeu, Auckland. Ph: 09 412 9817. www.adria.nz

Notice to Buyer: Careful tests have proved that the product is suitable for the recommended purposes when used in accordance with our instructions. However, since storage and use of product are beyond our control, we do not therefore accept any liability for damage or loss of efficacy which may result from the use of this product, whether used in accordance with the directions or not. We are liable for a consistent quality of the product, but the risk of its storage and use is not borne by us.

RESISTANCE MANAGEMENT: EroMite is a **GROUP 10 INSECTICIDE** and is a mite growth inhibitor. Resistance to this insecticide could develop from excessive use. To minimise this risk use strictly in accordance with the label instructions and resistance management strategies that exist for any pest listed on the label. EroMite may be applied only ONCE in total per crop cycle. EroMite should be used in a programme that incorporates insecticides with different modes of action to that of **GROUP 10 INSECTICIDES**. Refer to the NZPPS website (www.nzpps.org/resistance/index.php) for more detailed information.

