

GENERAL INFORMATION:

Evito is an effective fungicide for the control of primary and secondary powdery mildew on apples, field grown pumpkins and winter squash. Evito has contact, translaminar and vapour phase modes of action, all of which contribute to its eradicator and protectant properties. Evito is specific in its action against powdery mildews.

IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS.

DIRECTIONS FOR USE:

CROP	DISEASE	REMARKS
Apples — all varieties	Powdery mildew	Use 40-50mL/100 litres of water as a dilute spray or adjust accordingly for concentrate spraying. Use the higher rate when powdery mildew infection is moderate to high. Commence at tight cluster and continue at 10-14 day intervals until new leaf growth ceases. Use the closer interval during rainy weather or during severe powdery mildew infection periods. Do not apply more than three applications in one season.
Field grown pumpkins and winter squash	Powdery mildew	Use 800mL — 1 litre/ha at 10 to 14 day intervals. Use the higher rate and/or closer interval during periods of intense disease pressure. Use as a preventative application and ensure uniform coverage. Do not apply more than three applications in one season.

IT IS AN OFFENCE FOR USERS OF THIS PRODUCT TO CAUSE RESIDUES EXCEEDING THE RELEVANT MRL IN THE NEW ZEALAND FOOD NOTICE: MAXIMUM RESIDUE LEVELS FOR AGRICULTURAL COMPOUNDS.

Withholding period: Apples, field grown pumpkins and winter squash — 14 days. For apples grown for export consult your exporter for withholding period information.

Application: May be applied through all conventional orchard and boom spraying equipment. Maintain agitation during mixing and spray application.

Compatibility: Evito may be tankmixed with most commonly used insecticides, fungicides and calcium nitrate. Do not mix with very alkaline materials.

Rainfall: Rain within 6 hours following application may reduce effectiveness.

IPM: Evito is compatible with integrated pest management programmes based on natural predators and parasites.

HARMFUL

Keep out of reach of children

Evito®

For the control of powdery mildew on apples, field grown pumpkins and winter squash.

Active Ingredient: Contains 250g/L bupirimate in the form of an oil-in-water emulsion.

GROUP 8 FUNGICIDE

Registered pursuant to the ACVM Act 1997 No. P8300
See www.foodsafety.govt.nz for registration conditions
HSNO Classifications: 6.1E, 6.3B, 6.4A, 6.5B, 6.9B, 9.1A
EPA Approval Code: HSR100490

This substance must be under the personal control of an approved handler when the substance is applied in a wide dispersive manner or used by a commercial contractor.

In a transport emergency dial 111, Police or Fire Brigade.
For specialist advice in a First Aid emergency only call:
National Poisons Centre on 0800 764 766.



9 421903 419378 >

Net Contents: 5 Litres



WARNING: This substance may be harmful if swallowed, inhaled or absorbed through the skin. May cause internal organ damage from repeated oral exposure at high doses. May cause skin and eye irritation. May cause skin sensitisation from prolonged skin contact. Avoid skin contact or inhalation. Avoid eye contact.

APPROVED HANDLER: Required.

HANDLING PRECAUTIONS: EQUIPMENT: Apply using calibrated and maintained equipment in accordance with the NZ Standard for the Management of Agrichemicals (NZS8409). Must be applied via ground-based methods only.

STORAGE: Store in original container tightly closed and in a locked, dry, cool, well ventilated area, away from foodstuffs. Storage must be generally in accordance with the NZ Standard for the Management of Agrichemicals (NZS8409).

PRECAUTIONS: Avoid contact with skin and eyes, and inhalation of spray mist. Wash hands and exposed skin before meals and after work. Avoid contamination of any water supply with chemical or empty container. Apply this product carefully. Ensure compliance with relevant regional air management plans.

EVITO SHALL NOT BE APPLIED ONTO OR INTO WATER.

EVITO IS TO BE APPLIED VIA GROUND BASED APPLICATION SYSTEMS ONLY.

VERY TOXIC TO AQUATIC ORGANISMS. A strategy to minimise spray drift should be employed at all times when applying spray near aquatic environments.

DISPOSAL OF CONTAINER: Triple rinse container and add residue to spray tank. Return clean, empty container to an AgRecovery site for disposal. Avoid contamination of any water supply with chemical or empty container.



FIRST AID: If swallowed do NOT induce vomiting. For advice, contact the National Poisons Centre on 0800 POISON (0800 764766) or a doctor immediately. If spilt on the skin, remove contaminated clothing and wash affected areas of skin. DO NOT SCRUB THE SKIN. If concentrate is splashed in eyes flush with running water for at least 15 minutes. Take to hospital without delay.

SPILLS: Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely (such as to a suitable landfill).

Batch No. and Manufacturing Date: See on packaging.

Registered to and Distributed by: Adria New Zealand Limited.
P.O. Box 535, 407 State Highway 16 Kumeu, Auckland. Ph: 09 412 9817. Fax: 09 412 9807. www.adria.nz

Notice to Buyer: Careful tests have proved that the product is suitable for the recommended purposes when used in accordance with our instructions. However, since storage and use of product are beyond our control, we do not therefore accept any liability for damage or loss of efficacy which may result from the use of this product, whether used in accordance with the directions or not. We are liable for a consistent quality of the product, but the risk of its storage and use is not borne by us.

