



# Paraquat 200SL

A non-selective contact herbicide for the control of many annual and perennial grasses and broadleaf weeds in clover seed crops, drains, forestry, industrial sites, Lucerne, waterways and for the control of Barley grass.

---

<u>Active Ingredient:</u>	200g/litre paraquat as the dichloride salt
<u>Chemical Group:</u>	Bipyridyl
<u>Formulation:</u>	Soluble concentrate
<u>Pack Size:</u>	10 Litre
<u>Compatibility:</u>	Paraquat 200SL is compatible with most commonly used herbicides. As formulations of other manufacturer's products are beyond the control of Adria all mixtures should be tested prior to use.

---

**DANGER: KEEP OUT OF REACH OF CHILDREN. PARAQUAT CAN KILL IF SWALLOWED. NEVER PUT IN A DRINKS BOTTLE.**

**BEWARE:** Apply this product carefully. Spray drift may cause serious damage to other desirable plants. A strategy to minimize spray drift should be employed at all times when applying spray near aquatic environments.

**RESISTANT WEEDS WARNING:** Paraquat 200SL is a GROUP D HERBICIDE, with a Photosystem-I-electron diversion mode of action. Naturally occurring weed biotypes resistant to GROUP D herbicides can become more prevalent after many years of continuous use. To delay the onset of, or control resistant weeds, use in rotation with herbicides of different mode of action. Since the occurrence of resistant weeds is difficult to detect prior to use Adria New Zealand Ltd accepts no liability for losses that may result from the failure of this product to control resistant weeds.

**Before opening, carefully read warning statement, precaution instructions, first aid instructions and directions for use.**

Paraquat 200SL is inactivated by soil so plants should be clean of mud or dust at time of application and only clean water should be used. Paraquat 200SL is non-residual and translocated, so foliage of plants should have complete coverage of spray solutions to achieve best results.

**APPLICATION: BOOMSPrAY:** Use water volumes up to 300 litres/ha to ensure complete plant coverage. **KNAPSACK, HANDGUN:** To ensure complete coverage of plants use equipment and pressure that does not induce misting.

**MIXING:** Half fill the spray tank with water then add Paraquat 200SL. Agitate thoroughly while filling tank with water. When mixing with other herbicides, add Paraquat 200SL after other water soluble and after flowable products.

**RAINFALL:** Paraquat 200SL is rainfast within 20 minutes of drying on the plant.

**WEEDS CONTROLLED:** Most annual and perennial grasses except couch, paspalum and kikuyu. Clumpy grasses such as cocksfoot and tall fescue may have regrowth and require further treatment. Most annual broadleaf weeds are controlled when young and actively growing, however regrowth will occur with perennial broadleaf weeds.

**DIRECTIONS FOR USE:**

METHOD	SITUATION	RATE	CRITICAL COMMENTS
Handgun or knapsack	Barley grass control	300ml/100 litres of water	Spray when barley grass is at the green seedhead stage.
Boom	Clover Seed Crops. (Grass suppression only)	1.5-2.0 litres/ha	Application should be made 2 weeks prior to closing the paddock. Irrigate where appropriate 1-2 weeks post application.
Handgun Boom	Drainage Banks	750ml/100 litres of water 7.5 litres/ha	Spray weeds emerging from water only.
Boom	Lucerne Crops	2.0-3.0 litres/ha	Spray when lucerne is dormant and weeds are young and actively growing. For residual control add atrazine or simazine or other appropriate herbicides.
Boom	Pasture Renovation	1.5 – 2.0 litres/ha	Spray Autumn or Spring prior to reseedling. When there are high levels of perennial weeds, better results will be obtained using glyphosate.
As a directed spray	Strawberries, Vegetables, Tobacco (inter row only)	3.0 litres/ha	Use a low pressure nozzle with a shielded spray on weeds up to 7cm in height.
As a directed spray.	Citrus or deciduous fruit orchards. Forest planting and vineyards	350ml/100 litres of water	Apply to young actively growing weeds. Spray should be directed at the base of trees or vines. Avoid contact with green bark or foliage. Tank mix with a suitable residual herbicide if required.

**IT IS AN OFFENCE FOR USERS OF THIS PRODUCT TO CAUSE RESIDUES EXCEEDING THE RELEVANT MRL IN THE NEW ZEALAND (MAXIMUM RESIDUE LIMITS OF AGRICULTURAL COMPOUNDS) FOOD STANDARDS.**

Registered to and Distributed by: Adria New Zealand Limited. P.O. Box 535 Kumeu, Auckland 0841, NEW ZEALAND.  
Ph: +64-9-412-9817; Fax: +64-9-412-9807.

[www.adriacp.co.nz](http://www.adriacp.co.nz)

Paraquat 200SL is Registered pursuant to the ACVM Act 1997, Approval No. P8472. EPA Approval Code: HSR000828.