

Providor®

A fungicide for the control of Botrytis in grapes, boysenberries and strawberries.



Active Ingredient: 500g/litre fenhexamid

Chemical Group: Hydroxylanilide

Formulation: Suspension concentrate

Pack Size: 5L

Providor® offers highly effective Botrytis control:

- ✓ Long-lasting protection.
- ✓ IPM compatible.
- ✓ Key alternative fungicide for resistance management.
- ✓ No effect on grape fermentation.
- ✓ Rainfast in 1 hour.

About Providor®

Providor® fungicide provides excellent protection against botrytis infections during the crucial flowering window and beyond. Providor® is an ideal rotation partner with other specialised botryticides and has no known cross-resistance with other chemicals. Providor is soft on beneficials and compatible with most IPM programmes.

General Information

Providor® is a contact fungicide for the control of Botrytis in grapes, boysenberries and strawberries.

IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS.

Directions for use:

According to law, the listed restrictions on the use of this product must be observed, and the product must only be used according to the label instructions.

The recommended rates are for high volume spraying to the point of run-off.

For concentrate spraying adjust dilution rate accordingly.

Crop	Disease	Rate	Critical Comments
Grapes	Botrytis	75ml per 100 litres or a minimum of 750ml/ha.	Make two applications at pre-bunch closure and veraison or at veraison and again 14 days later. Apply in sufficient water to achieve good coverage of the bunches.
Boysenberries, Strawberries	Botrytis	150ml per 100 litres or a minimum of 1.5L/ha.	Make two applications over the flowering period ensuring good coverage of the flowers.

Mixing: Add the required amount of PROVIDOR to the partly filled spray tank, then add the remainder of the water while thoroughly agitating. Continue agitation while spraying.

Compatibility: PROVIDOR is compatible with most commonly used fungicides and insecticides. Care should be taken with mixtures with more than one chemical. All mixtures should be tested prior to use.

IT IS AN OFFENCE FOR USERS OF THIS PRODUCT TO CAUSE RESIDUES EXCEEDING THE RELEVANT MRL IN THE FOOD NOTICE: MAXIMUM RESIDUE LEVELS FOR AGRICULTURAL COMPOUNDS.

Withholding Periods:

Grapes – 21 days

Boysenberries – 35 days

Strawberries – 1 day

Note: Export crops: – contact your exporter, industry body or industry crop adviser.

Application: May be applied through all conventional, ground-based boom spraying equipment. Maintain agitation during mixing and spray application.