

IT IS AN OFFENCE FOR USERS OF THIS PRODUCT TO CAUSE RESIDUES EXCEEDING THE RELEVANT MRL IN THE NEW ZEALAND FOOD NOTICE: MAXIMUM RESIDUE LEVELS FOR AGRICULTURAL COMPOUNDS.

Directions for use:

CROP	WEEDS CONTROLLED	RATE	CRITICAL COMMENTS
Peas Process, Garden, Field	Annual poa, Black nightshade, Calandrinia, Chickweed, Annual mouse-ear chickweed, Clovers (seedlings), Cornbind, Fathen, Field pansy, Fumitory, Hedge mustard, Henbit, Redroot, Scarlet pimpernel, Scentless chamomile, Shepherd's purse, Speedwells, Spurrey, Storksbill, Twincross, Wild turnip, Willow weed, Wireweed. For best results, annual broadleaf weeds should be not greater than 4 true leaves, as larger weeds, especially Storksbill and Wireweed, are less susceptible. Note: Californian thistle and other perennial broadleaf weeds are not controlled.	0.75-1 litre in a minimum of 250 L of water/ha	As peas are a tender crop which depend on the waxy leaf surface for selectivity, Terbo Flo herbicide must be applied to actively growing pea crops less than 15cm high, not suffering from disease, wind damage, drought stress or checked by unseasonal cold, wet conditions. Use the crystal violet dye retention test to confirm the presence of sufficient leaf wax before spray application. After rain allow 2-3 days drying before application. Apply when the crop has 3-5 nodes (5-10cm high), using the lower rate when weeds are very small. Terbo Flo provides quick knockdown plus residual weed control. Best results will be obtained when applied to a fine moist seedbed. Weed control may be less effective if application is made when weeds are in a dry hardened growth condition. See compatibility statement regarding additives.
Potatoes	Use the lower rate where scarifying for weeds is carried out before crop canopy closure. Apply after moulting, free from clods and when the weeds are still small. Do not apply after 5% of the crop has emerged as damage will result.	1-2 litres in a minimum of 350 litres of water/ha	

RAINFALL: Providing the spray has dried on the weed foliage, rain following application will not impair results.

COMPATIBILITY: Do not mix with oils, surfactants or emulsifiable concentrate formulations or crop injury may result.

APPLICATION: Before using Terbo Flo, it is important that equipment is completely cleaned of all traces of oil, surfactants and emulsifiable concentrates. Apply only with conventional ground spraying equipment. Care should be taken to avoid excess overlapping. Maintain agitation throughout spraying. Do not use strainers finer than 300 microns.

Sprayer clean up: To avoid injury to subsequently sprayed crops, thoroughly remove all traces of Terbo Flo from mixing and spray equipment. Thoroughly clean sprayer with several flushes of clean water. Do not dispose of surplus spray liquid or washings on cropping land or in streams.

RESISTANT WEEDS WARNING: Naturally occurring weed biotypes resistant to triazine herbicides can become more prevalent after many years of continuous use. To delay the onset of, or control resistant weeds, use in rotation with herbicides of different mode of action. Since the occurrence of resistant weeds is difficult to detect prior to use Adria New Zealand Ltd accepts no liability for losses that may result from the failure of this product to control resistant weeds.

WARNING
KEEP OUT OF REACH OF CHILDREN

Terbo[®]FLO

For selective post-emergent control of annual broadleaf weeds in peas and potatoes.

Active Ingredient: contains 350 g/litre terbutryn plus 150 g/litre terbuthylazine in the form of a suspension concentrate.

GROUP C1 HERBICIDE

Registered pursuant to the ACVM Act 1997 No. P8195

See www.foodsafety.govt.nz for registration conditions

HSNO Classifications: 6.1D, 6.7B, 6.9B, 9.1A, 9.2B, 9.3C

EPA APPROVAL CODE: HSR100520

THIS PRODUCT MUST BE UNDER THE CARE OF AN APPROVED HANDLER DURING WIDE DISPERSIVE USE OR WHEN USED BY A COMMERCIAL CONTRACTOR.

In a transport emergency dial 111, Police or Fire Brigade.

For specialist advice in a First Aid emergency only call:

National Poisons Centre on 0800 764 766.



Net Contents: 5 litres



WARNING: This substance may be harmful if swallowed, inhaled or absorbed through the skin, and may cause irritation to eyes. Avoid eye contact. Avoid inhalation of vapours or mist from spray. May cause skin irritation. Avoid skin contact. May cause internal organ damage from repeated oral exposure at high doses. Suspected carcinogen.

PRECAUTIONS: Store in original container, tightly closed, away from foodstuffs in a cool dry, secured area. Avoid contact with skin and eyes, and inhalation of spray mist. When mixing or applying wear appropriate clothing including impervious, elbow-length gloves and eye protection. Wash hands and exposed skin before meals and after work. Avoid contamination of any water supply with chemical or empty container.

Apply this product carefully. Spray drift may cause serious damage to desirable plants. Ensure compliance with relevant regional air management plans.

VERY TOXIC TO AQUATIC LIFE WITH LONG-LASTING EFFECTS. A strategy to minimise spray drift should be employed at all times when applying spray near aquatic environments.

MUST NOT BE APPLIED ONTO OR INTO WATER.

VERY TOXIC IN THE SOIL ENVIRONMENT. HARMFUL TO TERRESTRIAL VERTEBRATES.

DISPOSAL OF CONTAINER: Triple rinse container and add residue to spray tank. Return clean, empty container to an AgRecovery site for disposal. Avoid contamination of any water supply with chemical or empty container.



FIRST AID: If swallowed do NOT induce vomiting. For advice, contact the National Poisons Centre on 0800 POISON (0800 764766) or a doctor immediately. If spilt on the skin, remove contaminated clothing and wash affected areas of skin. DO NOT SCRUB THE SKIN. If concentrate is splashed in eyes flush with running water for at least 15 minutes. Take to hospital without delay.

SPILLS: Wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash area with water and absorb with further inert material. Dispose of waste safely (such as to a suitable landfill).

Batch No. and Manufacturing Date: See on packaging.

Registered to and Distributed by:

Adria New Zealand Limited.

P.O. Box 535, 407 State Highway 16 Kumeu, Auckland.

Ph: 09 412 9817. Fax: 09 412 9807. www.adria.nz



Notice to Buyer: Careful tests have proved that the product is suitable for the recommended purposes when used in accordance with our instructions. However, since storage and use of product are beyond our control, we do not therefore accept any liability for damage or loss of efficacy which may result from the use of this product, whether used in accordance with the directions or not. We are liable for a consistent quality of the product, but the risk of its storage and use is not borne by us.

BEWARE: Apply this product carefully. Spray drift may cause serious damage to other desirable plants. A strategy to minimize spray drift should be employed at all times when applying spray near aquatic environments.

GENERAL INSTRUCTIONS

Terbo FLO is absorbed by foliage and roots to ensure quick knockdown. It translocates from roots to foliage and inhibits photosynthesis.